

Institut für
Biologische Analytik
und Consulting GmbH

Study Report

Laboratory testing for toxicity (acute contact and oral LD₅₀) of Confidor SC 200 to honey bees (*Apis mellifera* L.) (Hymenoptera, Apidae)

(GLP compliant study according to EPPO 170 (1992))

Author: Dipl. Bio. [REDACTED]

Study completion date: September 9, 1995

Study performed for Sponsor:

Bayer AG
Pflanzenschutz
Ökobiologie
D-51368 Leverkusen

Testing facility


IBACON
Institut für
Biologische Analytik
und Consulting GmbH
D - 64380 Rossdorf

Project 790036



790036 / MO-00-007453

Copy of GLP Certificate



Dit document is geen eigen
Op dit document
Publicatie, verspreiding
recht

Consequently, w

of its contents

Contents

Copy of GLP Certificate..... 2

List of tables and figures 4

1. Summary 5

2. Survey of the study..... 6

 2.1 General 6

 2.2 Quality assurance 7

 2.3 Archiving..... 7

 2.4 Signatures of project staff 8

3. Quality Assurance Unit 9

4. Statement of compliance 10

5. Objectives of study 11

 5.1 Introduction 11

6. Guidelines..... 11

7. Material and Methods..... 12

 7.1 Test substance and positive control substance 12

 7.2 Test system 13

 7.3 Test units 13

 7.4 Test conditions 13

 7.5 Food..... 13

 7.6 Application of the test substance and of the controls..... 13

 7.7 Course of the test..... 14

 7.7.1 Contact toxicity test..... 14

 7.7.2 Oral toxicity test 14

 7.8 Test parameter 15

 7.9 Evaluation of the results..... 15

 7.10 Deviations from the protocol..... 15

8. Results and discussion..... 17

 8.1 Contact toxicity test..... 17

 8.2 Oral toxicity test..... 20

 8.3 Conclusions 22

9. Literature 23

10. Distribution of the report..... 23

Appendix 24

List of tables

Summary table 5

Table 1: Mortality and Behavioural abnormalities in the contact toxicity test 18

Table 2: Mortality and behavioural abnormalities in the oral toxicity test 20

Table 3: Mortality and behavioural abnormalities of the bees under the influence of several dosages of the test substance in course of the contact toxicity test24

Table 4: Mortality and behavioural abnormalities of the bees under the influence of several dosages of the test substance in the oral toxicity test 26

Table 5: Oral intake test: weight of syringes 28

List of figures

Figure 1: Mortality and LD₅₀ in the contact test 19

Figure 2: Mortality and LD₅₀ in the oral test 21

Dit document is geen eigendom van het Ctgb en wordt beschikbaar gemaakt op grond van een wettelijke verplichting tot openbaarmaking. Op dit document kunnen rechten van derden rusten, waaronder intellectuele eigendomsrechten en/of auteursrechten. Voorts kan dit document onder een regeling omtrent gegevensbescherming vallen. Het gebruik van dit document of de inhoud hiervan zonder de toestemming van de rechthebbende van dit document kan derhalve verboden zijn en een inbreuk opleveren van de rechten van deze rechthebbende.

This document is not the property of the Ctgb and only provided based on mandatory freedom of information requirements. The document may be subject to rights such as intellectual property and copy rights of third parties. Furthermore, this document may fall under a regulatory data protection regime. Consequently, any publication, distribution, reproduction and/or publishing and any commercial exploitation and use of this document or its contents without the permission of the owner of this document may therefore be prohibited and violate the rights of its owner.

1. Summary

The contact and oral LD₅₀ (24 h and 48 h) of Confidor SC 200 to honey bees were tested according to EPPO 170 (1992) and GLP regulations. Confidor SC 200 was applied in five dosages (contact and oral toxicity test), one solvent control, one untreated negative control (contact test) and one positive control with toxic standard (Dimethoate 0.2 µg a.i./bee). The following dosages of the test substance were tested in three replicates of ten bees each:

Contact toxicity test			Oral toxicity test		
Test Substance Dosage (nominal) µg / bee	Mortality		Test Substance Dosage µg / bee	Mortality	
	24 h (%)	48 h (%)		24 h (%)	48 h (%)
5.0	70.0	86.7	0.169	40.0	80.0
1.0	60.0	86.7	0.083	20.0	23.3
0.5	50.0	66.7	0.0173	6.7	10.0
0.1	16.7	16.7	0.0083	0.0	0.0
0.05	20.0	20.0	0.0018	0.0	0.0
calculated LD ₅₀ µg/bee (95 % confidence range)	0.73 (0.38 to 1.39)	0.29 (0.19 to 0.45)	calculated LD ₅₀ µg/bee (95 % confidence range)	> 0.169	0.103 (0.073 to 0.144)
Controls	Mortality		Controls	Mortality	
	24 h (%)	48 h (%)		24 h (%)	48 h (%)
untreated Control	0.0	0.0	-	-	-
solvent control	0.0	0.0	negative control	0.0	3.3
positive control	73.3	73.3	positive control	96.7	100.0

The results of this study show toxic effects of Confidor SC 200 to honey bees in the contact and oral toxicity test. The acute contact LD₅₀ (48h) was calculated to be 0.29 µg/bee and the acute oral LD₅₀ (48h) was calculated to be 0.103 µg/bee.

2. Survey of the study

2.1 General

Title

Laboratory testing for toxicity (acute contact and oral LD₅₀) of
Confidor SC 200 to honey bees (*Apis mellifera* L.)
(Hymenoptera, Apidae)

Sponsor:

Bayer AG
Pflanzenschutz
Ökobiologie
D-51368 Leverkusen

Monitoring:

[REDACTED]

Test substance:

Confidor SC 200

Testing facility:

IBACON
Institut für Biologische Analytik
und Consulting GmbH
Industriestrasse 1
D-64380 Rossdorf

IBACON Projectnumber: 790036

Project staff

Management: [REDACTED]

Study director: Dipl. Biol. [REDACTED]

Quality Assurance Unit: [REDACTED]

Schedule

Date of protocol and German translation:	May 11, 1995
Date of 1st amendment to protocol:	June 29, 1995
Experimental start date:	July 24, 1995
Experimental termination date:	July 27, 1995
Date of draft report:	August 10, 1995
Date of report:	September 9, 1995

2.2 Quality assurance

This study was performed in compliance with:

- 'The OECD principles of Good Laboratory Practice' (Paris, France 1992)
- Chemikaliengesetz (Chemicals Act) of the Federal Republic of Germany, Anlage 1 (Annex 1), dated July 25, 1994 (BGBL I 1994, p. 1703).

This study was assessed to assure compliance with the protocol and the IBACON Standard Operating Procedures. The study and/or testing facility were periodically inspected by the Quality Assurance Unit (QAU) and the dates and phases of the inspections are included with the study report. The data contained with the report were audited. A quality assurance statement, signed by the QAU-Manager, is included in the report.

2.3 Archiving

IBACON, Institut für Biologische Analytik und Consulting GmbH, Industriestrasse 1, D 64380 Rossdorf, will archive the following data for 30 years: - all raw data, the protocol and a certified copy of the final report.

The following sample will be archived for at least 2 years following the date on which the report is audited by the Quality Assurance Unit: - sample of the test article.

No raw data or material relating to the study will be discarded without the sponsor's prior consent.

Dit document is geen eigendom van het Ctgb en wordt beschikbaar gemaakt op grond van een wettelijke verplichting tot openbaarmaking. Op dit document kunnen rechten van derden rusten, waaronder intellectuele eigendomsrechten en/of auteursrechten. Voorts kan dit document onder een regeling omtrent het gebruik van afgedrukt document of de inhoud hiervan vande toestemming vande rechthebbende van dit document kan derhalve verboden zijn en een inbreuk opleveren van de rechten van de rechthebbende.

This document is not the property of the Ctgb and only provided based on mandatory freedom of information requirements. The document may be subject to rights such as intellectual property and copy rights of third parties. Furthermore, this document may fall under a regulatory data protection and any commercial exploitation and use of this document or its contents without the permission of the owner of this document may therefore be prohibited and violate the rights of its owner.

2.4 Signatures of project staff

Study Director:

Dipl. Biol. [redacted]

[redacted signature]

date:

September 11, 1995

Management:


[redacted signature]

[redacted signature]

date:

September 11, 1995

3. Quality Assurance Unit




Dit docu
Pu

Conse

nts

4. Statement of compliance



Dit document is niet te verspreiden of openbaar te maken. Het is vertrouwelijk en kan wettelijk beschermd zijn. Het verspreiden of openbaar maken van dit document is strafbaar. Het verspreiden of openbaar maken van dit document is strafbaar. Het verspreiden of openbaar maken van dit document is strafbaar.

Publicatie, verspreiden of openbaar maken van dit document is strafbaar. Het verspreiden of openbaar maken van dit document is strafbaar. Het verspreiden of openbaar maken van dit document is strafbaar.

This document is not to be disseminated or made public. It is confidential and may be legally protected. The dissemination or public release of this document is punishable. The dissemination or public release of this document is punishable. The dissemination or public release of this document is punishable.

Consequently, any publication, distribution or public release of this document without the permission of the author is prohibited. Consequently, any publication, distribution or public release of this document without the permission of the author is prohibited. Consequently, any publication, distribution or public release of this document without the permission of the author is prohibited.

er.
or its contents

5. Objectives of study

5.1 Introduction

Honey bees (*A. mellifera*) can be impacted by plant protection products. If the proposed use pattern of Confidor SC 200 indicates a possible exposure of honey bees, this study shall be useful for the registration of the plant protection products in question.

6. Guidelines

This study was based on the procedures indicated by the following internationally accepted guidelines:

EPPO 1992: Guideline on test methods for evaluating the side-effects of pesticides on honey bees,

Bulletin OEPP/EPPO Bulletin 22, 203-215 1992, No. 170

Dit document is geen eigendom van het Ctgb en wordt beschikbaar gemaakt op grond van een wettelijke verplichting tot openbaarmaking.
Op dit document kunnen rechten van derden rusten, waaronder intellectuele eigendomsrechten en/of auteursrechten.
Voorts kan dit document onder een regeling omtrent gegevensbescherming vallen.
Publicatie, verspreiding, vermenigvuldiging, commerciële exploitatie en gebruik van dit document of de inhoud hiervan zonder de toestemming van de rechthebbende van dit document kan derhalve verboden zijn en een inbreuk opleveren van de rechten van deze rechthebbende.

This document is not the property of the Ctgb and only provided based on mandatory freedom of information requirements.
The document may be subject to rights such as intellectual property and copy rights of third parties.
Furthermore, this document may fall under a regulatory data protection regime.
Consequently, any publication, distribution and/or publishing and any commercial exploitation and use of this document or its contents without the permission of the owner of this document may therefore be prohibited and violate the rights of its owner.

7. Material and methods

7.1 Test substance and positive control substance

Name: Confidor SC 200
Formulation - No.: 0458 (according to 03554/0456)
Tox-No.: 3865-00
Product-No.: 00173846
Active ingredients/Purity: 205.9 g/l Imidacloprid
Certificate of Analysis date: April 19, 1995
Use: Insecticide
Expiration date: October 13, 1995
Aggregate State at RT: liquid (suspension)

Colour: white
Density: 1.101 g/ml
Solubility in: water: suspensible (dispersible)

Stability:
Hydrolysis of the active ingredient: DT₅₀ of a.i. > 1 year (pH 7)
DT₅₀ of a.i. ca. 1 year (pH 9)

Sample received: May 5, 1995; 12.00 a.m.
Storage: at room temperature

Positive control substance

Name: Perfekthion
Active Ingredient: 400 g/l (37 %) Dimethoate
Manufacturer: BASF AG, D-67056 Ludwigshafen
Batch - No.: 98 0878

Applied Amount in this Study: 0.5 µl of formulation (corresponding to 0.2 µg active ingredient) per bee

Expiration Date: 11/1995
Storage: in original container, protected from light at aprox. 20°C.

Safety Precautions: routine hygienic procedures will be sufficient to assure personnel health and safety.

7.2 Test system

The study was performed with worker honey bees (*Apis mellifera* L.), about four to six weeks old, bred by IBACON in normal beekeeper's manner (responsible beekeepers: [REDACTED], Dipl. Biol. [REDACTED]). For the tests, the bees were caught randomly from the entrance hole and from the edge positioned honeycombs of the hives in groups of ten with glass capture tubes. The capture took place without anaesthetics ultimately before the experiments started.

7.3 Test units

Stainless steel chambers (width 10 cm, height 8.5 cm, depth 5.5 cm) served as test cages. The front side of the cages was closed with removable glass sheets; the bottom is perforated with ventilation holes (\varnothing 1 mm). The top contains two openings for the food tube and entry for the bees. The inner sides of the cages (except front side) were covered with filter paper (Co. Machery & Nagel, D-52355 Düren, Art.No. 68). These test cages allowed an adequate rearing of the bees during the experiment combined with an easy observation through the front glass slide.

7.4 Test conditions

The test cages were exposed in incubators at 27 - 29 °C and 45 - 80 % relative humidity and darkness instead of 25 ± 2 °C and 60 - 70 % relative humidity as mentioned in the EPPO guideline No. 170. It can be excluded that this change had an influence on the results of this study (see 7.10 Deviation to the protocol).

Temperature and relative humidity were continuously recorded with a Thermo-Hygrometer. The incubators were ventilated to avoid possible accumulations of pesticide vapour.

7.5 Food

During the tests, the bees were provided *ad libitum* with commercial ready to use syrup for honey bees as food (Apiinvert, Co. Südzucker AG, D-97199 Ochsenfurt).

7.6 Application of the test substance and of the controls

Each test was performed in five dosages (acute contact and oral toxicity test) of Confidor SC 200, one solvent control, one untreated negative control and one positive control with toxic standard (Dimethoate) with three replicates per dosage / control. One replicate was formed by one test cage containing 10 honey bees.

7.7 Course of the test

7.7.1 Contact toxicity test

Five dosages of the test substance were tested in order to provide a rational base for a proper assessment of the contact LD₅₀ of Confidor SC 200 to honey bees.

Five solutions of the test substance in acetone/water (=solvent; 1:1 v/v) were prepared in adequate concentrations to yield the corresponding amounts per bee when 5 µl of the solution per individual were applied. The stock solution was prepared using 50 mg Test substance suspended in 50 ml solvent (highest dosage group; 1 µg/µl corresponding to 5 µg/bee/5 µl). This stock solution was diluted down to 0.01 µg/µl in four steps. The following dosages were tested: 5; 1; 0.5; 0.1 and 0.05 µg/bee.

For the negative control the bees only were anaesthetized. For the solvent-control 5 µl/bee of the pure solvent was used.

For the positive control 0.2 µg/bee dimethoate in 5 µl solvent was used.

Immediately after preparing the test substance dilutions the application was done. Ultimately before applying the test substance the bees were anaesthetized with CO₂. The CO₂ was given for 20 seconds directly into the test cages using one of the top openings. The anaesthetized bees were laid, ventral surface up, on filter paper in petri dishes. One 5.0 µl drop per bee of Confidor SC 200 in solvent was placed on the ventral thorax using a Burkard-Applicator. The treated bees were then returned into the test cages and kept under test conditions for 48 hours. The duration of the anaesthetization was between 140 and 300 seconds until the total recovery of all individuals.

7.7.2 Oral toxicity test

Five dosages of the test substance, one control in water (solvent/ready to use syrup) and one positive control in ready to use syrup with dimethoate were tested in order to provide a rational base for a proper assessment of the oral LD₅₀ of Confidor SC 200 to honey bees.

Five dilutions of the test substance in solvent/ready to use syrup (1 part solvent + 19 parts syrup) were prepared. 19.9 mg test substance was suspended in 200 ml water (= stock solution; corresponding to 99.5 µg/ml). 500 µl of this suspension was mixed with 9.5 ml Apiinvert (highest dosage group; ca. 0.1 µg/bee). The stock solution was diluted in 4 steps to yield the corresponding amounts per bee when ca. 30 mg of the dilution per individual were administered.

Test cages containing 10 bees each were prepared without food, letting the bees starve for 70 - 80 minutes. Within starving time of the bees preparation of the test substance dilutions was done. Following this, approximately 330 mg of the prepared dilutions in 1-ml syringes were hung into each cage through one of the top openings. The bees were observed as long as uptake of the dilution took place (uptake of contaminated food lasted not more than one hour). After the complete uptake of the test substance dilution the bees were provided with normal food (see 7.5).

The exact amount of test dilution consumed was determined by reweighing the syringes.

According to the weight results the following exact dosages were tested (details are given in table 5):

- 0.169 µg / bee
- 0.083 µg / bee
- 0.0173 µg / bee
- 0.0083 µg / bee
- 0.0018 µg / bee
- solvent/pure ready to use syrup control
- positive control (0.33 µg dimethoate / bee)

7.8 Test parameter

The experiments were started in the early midday to ensure a long enough observation time during the first day. The experiments were finished after 48 hours. Observation of the bees was continuous during the first 30 minutes after application of the test substance and then took place after the following times:

- 45 minutes, 60 minutes, 2 hours, 4 hours (first day)
- 24 hours, 48 hours (following days).

Any mortality and / or poisoning or behavioural abnormalities of the bees (e.g. food refuse, apathy, moving coordination problems) were recorded.

7.9 Evaluation of the results

Toxicity assessment of the test substance took place as comparison of the results obtained from the bees treated with test substance to those results obtained from the controls.

The study is valid because the mortality of the bees in the controls did not exceed 10 % within 24 hours and 48 hours.

The contact and oral LD₅₀ (24 h and 48 h) of the test substance was estimated with probit analysis according to Finney 1971.

7.10 Deviations from the protocol

Test conditions:

Because of technical reasons the test containers were exposed at 48 - 85 % rel. humidity, instead of 40 - 70 % rel. humidity as described in the protocol and as mentioned in the EPPO guideline No. 170 (60 - 70 %).

Contact Toxicity Test:

The solvent was placed on the ventral thorax using a Burkard - Applicator, instead of a GC - syringe as described in the protocol, because using a Burkard - Applicator leads to an improvement of the study.

These deviations had no influence on the validity of the study.

*Dit document is geen eigendom van het Ctgb en wordt beschikbaar gemaakt op grond van een wettelijke verplichting tot openbaarmaking.
Op dit document kunnen rechten van derden rusten, waaronder intellectuele eigendomsrechten en/of auteursrechten.
Voorts kan dit document onder een regeling omtrent gegevensbescherming vallen.
Publicatie, verspreiding, vermenigvuldiging, commerciële exploitatie en gebruik van dit document of de inhoud hiervan zonder de toestemming van de rechthebbende van dit document kan derhalve verboden zijn en een inbreuk opleveren van de rechten van deze rechthebbende.*

*This document is not the property of the Ctgb and only provided based on mandatory freedom of information requirements.
The document may be subject to rights such as intellectual property and copy rights of third parties.
Consequently, any publication, distribution, reproduction and/or publishing and any commercial exploitation and use of this document or its contents without the permission of the owner of this document may therefore be prohibited and violate the rights of its owner.*

Table 1: Mortality and behavioural abnormalities of the bees in the contact toxicity test; presented results are averages from three replicates (10 bees each) per dosage/control.
See appendix, table 3 for details.

test article dosage µg/bee	first 30 minutes		after 1 hour		after 4 hours		after 24 hours		after 48 hours	
	mortality average %	behav. abnorm. average %	mortality average %	behav. abnorm. average %	mortality average %	behav. abnorm. average %	mortality average %	behav. abnorm. average %	mortality average %	behav. abnorm. average %
5.0	0.0	100.0	0.0	100.0	10.0	90.0	70.0	30.0	86.7	10.0
1.0	0.0	3.3	3.3	96.7	13.3	86.7	60.0	40.0	86.7	3.3
0.5	0.0	0.0	0.0	76.7	0.0	100.0	50.0	50.0	66.7	0.0
0.1	0.0	0.0	0.0	6.7	10.0	10.0	16.7	0.0	16.7	0.0
0.05	0.0	0.0	0.0	0.0	13.3	3.3	20.0	0.0	20.0	0.0
untr.	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
solv.	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
posit.	0.0	13.3	0.0	0.0	26.7	26.7	73.3	0.0	73.3	0.0





Fig. 1: Mortality of the honey bees in course of the contact test; above: mortality and LD₅₀ within 24 hours; below: mortality and LD₅₀ within 48 hours;

8.2 Oral toxicity test

The amount of contaminated food ingested by the bees ranged from 29 to 38 mg/bee. According to the food uptake, the average dosages were calculated to 0.169, 0.083, 0.0173, 0.0083 and 0.0018 µg/bee (compare appendix table 5).

Ingestion of 0.169 µg/bee Confidor SC 200 resulted in 80 % mortality within 48 hours. The minimum mortality was 10 % (0.0173 µg/bee). No mortality was observed after the ingestion of 0.0083 and 0.0018 µg/bee.

100 % of the bees in the positive control group died within the experimental time (48 h). In the negative control group one bee (3.3 %) died within the whole experimental time.

The results of this experiment give a dose-effected relationship.

The mortality after 48 hours was 80 %, 23.3 % and 10 % in the 0.169 , 0.083 and 0.0173 µg/bee treatment groups.

According to these data, the acute oral LD₅₀ (24 h) was calculated to be > 0.169 µg/bee and the acute oral LD₅₀ (48 h) was calculated to be 0.103 µg/bee (95 % confidence range from 0.073 to 0.144 µg/bee; probit analysis).

Table 2: Mortality and behavioural abnormalities of the bees in the oral toxicity test; presented results are averages from three replicates (10 bees each) per dosage/control. negative = negative control, positive = positive control (treated with dimethoate). See appendix, table 4 for details.

uptaken test substance µg/bee	first 30 minutes		after 1 hour		after 4 hours		after 24 hours		after 48 hours	
	mortality average	behav. abnorm. average	mortality average	behav. abnorm. average	mortality average	behav. abnorm. average	mortality average	behav. abnorm. average	mortality average	behav. abnorm. average
	%	%	%	%	%	%	%	%	%	%
0.169	0.0	0.0	0.0	23.3	0.0	56.7	40.0	6.7	80.0	0.0
0.083	0.0	0.0	0.0	0.0	6.7	3.3	20.0	0.0	23.3	0.0
0.0173	0.0	0.0	0.0	0.0	6.7	0.0	6.7	0.0	10.0	0.0
0.0083	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
0.0018	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
negative	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3.3	0.0
positive	3.3	0.0	6.7	0.0	40.0	40.0	96.7	3.3	100.0	0.0





Fig. 2: Mortality of the honey bees in course of the oral test; above: mortality within 24 hours; below: mortality and LD₅₀ within 48 hours;

8.3 Conclusions

The results of this study exhibit effects of Confidor SC 200 to honey bees in the contact and oral toxicity test. The contact LD₅₀ of Confidor SC 200 was calculated to be 0.29 µg/bee (48 h) and 0.73 µg/bee (24 h) respectively.

The oral LD₅₀ of Confidor SC 200 was calculated to be >0.169 µg/bee (24 h) and 0.103 µg/bee (48 h) respectively.

*Dit document is geen eigendom van het Ctgb en wordt beschikbaar gemaakt op grond van een wettelijke verplichting tot openbaarmaking.
Op dit document kunnen rechten van derden rusten, waaronder intellectuele eigendomsrechten en/of auteursrechten.
Voorts kan dit document onder een regeling omtrent gegevensbescherming vallen.
Publicatie, verspreiding, vermenigvuldiging, commerciële exploitatie en gebruik van dit document of de inhoud hiervan zonder de toestemming van de rechthebbende van deze rechthebbende.*

*This document is not the property of the Ctgb and only provided based on mandatory freedom of information requirements.
The document may be subject to rights such as intellectual property and copy rights of third parties.
Furthermore, this document may fall under a regulatory data protection and use of this document or its contents
Consequently, any publication, distribution and/or publishing and any commercial exploitation and use of this document or its contents without the permission of the owner of this document may therefore be prohibited and violate the rights of its owner.*

9. Literature

EC Agrochemical Registration Directive (DS65) (Directive 91/414/EEC). ANNEX III. Requirements for the dossier to be submitted for the authorization of a plant production product

EPPO 1992: Guideline on test methods for evaluating the side-effects of plant protection products on honey bees, Bulletin OEPP/EPPO Bulletin 22, 203-215 1992, No. 170

Finney D.J. 1971: Probit Analysis. 3rd Edition, Cambridge University Press, London

10. Distribution of the report

Sponsor: 2x (original and one copy)

Study director: 1x (copy)

Dit document is geen eigendom van het Ctgb en wordt beschikbaar gemaakt op grond van een wettelijke verplichting tot openbaarmaking.
Op dit document kunnen rechten van derden rusten, waaronder intellectuele eigendomsrechten en/of auteursrechten.
Voorts kan dit document onder een regeling omtrent gegevensbescherming en/of de inhoud hiervan zonder de toestemming van de rechthebbende van dit document kunnen worden openbaar gemaakt.
Publicatie, verspreiding, vermenigvuldiging, commerciële exploitatie en gebruik van dit document of de rechten van deze rechthebbende, zonder de toestemming van de rechthebbende, is strafbaar.
Consequently, any publication, distribution, reproduction and/or publishing of this document may therefore be prohibited and violate the rights of its owner.
The document may be subject to rights such as intellectual property and copy rights of third parties.
Furthermore, this document may fall under a regulatory data protection regime.
Consequently, any publication, distribution, reproduction and/or publishing of this document may therefore be prohibited and violate the rights of its owner.

Appendix

Table 3: definitive contact toxicity test; mortality and behavioural abnormalities of the bees under the influence of several dosages of the test substance;

Observations: mortality and behavioural abnormalities; # = number of individuals; sympt. = observed symptoms according to the following key:

1 = vomit; 2 = moving coordination problems; 3 = apathy; 4 = intensive cleaning

vial	test article dosage	first 30 minutes			after 45 minutes			after 1 hour			after 2 hours			after 4 hours			after 24 hours			after 48 hours		
		dead	behav.abnor.		dead	behav.abnor.		dead	behav.abnor.		dead	behav.abnor.		dead	behav.abnor.		dead	behav.abnor.		dead	behav.abnor.	
#	µg/bee	#	#	sympt	#	#	sympt	#	#	sympt	#	#	sympt	#	#	sympt	#	#	sympt	#	#	sympt
1	5.0	0	10	1,2,3	0	10	1,2,3	0	10	1,2,3	0	10	2,3	1	9	3	6	4	1,2,3	9	1	2
2	5.0	0	10	1,2,3	0	10	1,3	0	10	1,2,3	0	10	2,3	0	10	3	7	3	1,2,3	7	2	2
3	5.0	0	10	1,2,3	0	10	1,3	0	10	1,2,3	0	10	2,3	2	8	3	8	2	1,2,3	10	0	0
1	1.0	0	1	1,3	1	7	1,2,3	1	9	1,2,3	1	9	2,3	3	7	1,2,3	4	6	1,2,3	6	1	3
2	1.0	0	0		0	0		0	10	2,3	0	8	2,3	1	9	2,3	8	2	1,2,3	10	0	0
3	1.0	0	0		0	0		0	10	2,3	0	10	2,3	0	10	1,2,3	4	4	1,2,3	10	0	0
1	0.5	0	0		0	1	3	0	3	3,2	0	3	2,3	0	10	2	5	0		5	0	0
2	0.5	0	0		0	5	2	0	10	1,2,3	0	2	2	0	10	2,3	6	4	2	10	0	0
3	0.5	0	0		0	6	2	0	10	1,2,3	0	4	2	0	10	2,3	4	6	2,3	5	0	0
1	0.1	0	0		0	0		0	1	3	0	1	2	1	0		1	0		1	0	0
2	0.1	0	0		0	0		0	1	2	0	1	2	0	0		1	0		1	0	0
3	0.1	0	0		0	0		0	0		0	0		2	3	1	3	0		3	0	0
1	0.05	0	0		0	0		0	0		0	0		0	0		0	0		0	0	0
2	0.05	0	0		0	0		0	0		0	0		0	1	2	2	0		2	0	0
3	0.05	0	0		0	0		0	0		0	3	3	4	0		4	0		4	0	0
1	untreated control	0	0		0	0		0	0		0	0		0	0		0	0		0	0	0
2	control	0	0		0	0		0	0		0	0		0	0		0	0		0	0	0
3	control	0	0		0	0		0	0		0	0		0	0		0	0		0	0	0
1	solvent control	0	0		0	0		0	0		0	0		0	0		0	0		0	0	0
2	control	0	0		0	0		0	0		0	0		0	0		0	0		0	0	0
3	control	0	0		0	0		0	0		0	0		0	0		0	0		0	0	0
1	positive control	0	2	2	0	5	1,2,4	0	0		0	8	2,3	7	2	2	10	0		10	0	0
2	control	0	2	2	0	0		0	0		0	0		0	6	2,3	6	0		6	0	0
3	control	0	0		0	0		0	0		0	0		1	8	2,3	6	0		6	0	0

Table 3 (continued): definitive contact toxicity test; mortality and behavioural abnormalities of the bees
average ± standard deviation from three replicates per concentration / control.

avg. ± sd = average ± standard deviation; behav. abnorm. = behavioural abnormalities;

test article dosage µg/bee	first 30 minutes				after 1 hour				after 4 hours				after 24 hours				after 48 hours			
	mortality		behav.abnor.		mortality		behav.abnor.		mortality		behav.abnor.		mortality		behav.abnor.		mortality		behav.abnor.	
	avg. ± sd		avg. ± sd		avg. ± sd		avg. ± sd		avg. ± sd		avg. ± sd		avg. ± sd		avg. ± sd		avg. ± sd		avg. ± sd	
	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%
5.0	0.0	0.0	100.0	0.0	0.0	0.0	100.0	0.0	10.0	10.0	90.0	10.0	70.0	10.0	30.0	10.0	86.7	15.3	10.0	10.0
1.0	0.0	0.0	3.3	5.8	3.3	5.8	96.7	5.8	13.3	15.3	86.7	15.3	60.0	20.0	40.0	20.0	86.7	23.1	3.3	5.8
0.5	0.0	0.0	0.0	0.0	0.0	0.0	76.7	40.4	0.0	0.0	100.0	0.0	50.0	10.0	50.0	10.0	86.7	28.9	0.0	0.0
0.10	0.0	0.0	0.0	0.0	0.0	0.0	6.7	5.8	10.0	10.0	10.0	17.3	16.7	11.5	0.0	0.0	16.7	11.5	0.0	0.0
0.05	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	13.3	23.1	3.3	5.8	20.0	20.0	0.0	0.0	20.0	20.0	0.0	0.0
untr.	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
solv.	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
posit.	0.0	0.0	13.3	11.5	0.0	0.0	0.0	0.0	26.7	37.9	26.7	37.9	73.3	23.1	0.0	23.1	73.3	23.1	0.0	0.0

Table 4: definitive oral toxicity test; mortality and behavioural abnormalities of the bees under the influence of several dosages of the test substance; test article dosages calculated after reweighing the syringes.

Observations: mortality and behavioural abnormalities; # = number of individuals; sympt. = observed symptoms according to the following key;

1 = food refuse/vomit; 2 = moving coordination problems; 3 = apathy; 4 = intensive cleaning

vial #	test article dosage µg/bee	first 30 minutes			after 45 minutes			after 1 hour			after 2 hours			after 4 hours			after 24 hours			after 48 hours		
		dead #	behav. abnor #	symp	dead #	behav. abnor #	symp	dead #	behav. abnor #	symp	dead #	behav. abnor #	symp	dead #	behav. abnor #	symp	dead #	behav. abnor #	symp	dead #	behav. abnor #	symp
1	0.172	0	0		0	1	2	0	1	2	0	9	3	0	4	3	4	0		5	0	
2	0.179	0	0		0	1	2	0	4	2	0	10	3	0	5	3	6	1	3	10	0	
3	0.158	0	0		0	0		0	2	2	0	10	3	0	8	3	2	1	3	9	0	
1	0.080	0	0		0	0		0	0		0	0	0	0	0	0	1	0		2	0	
2	0.089	0	0		0	0		0	0		0	0	0	0	0	0	2	0		2	0	
3	0.080	0	0		0	0		0	0		0	3	3	2	1	3	3	0		3	0	
1	0.017	0	0		0	0		0	0		0	1	3	1	0	0	2	0		2	0	
2	0.018	0	0		0	0		0	0		0	0	0	0	0	0	0	0		0	0	
3	0.017	0	0		0	0		0	0		0	1	2	1	0	0	0	0		1	0	
1	0.0073	0	0		0	0		0	0		0	0	0	0	0	0	0	0		0	0	
2	0.0086	0	0		0	0		0	0		0	0	0	0	0	0	0	0		0	0	
3	0.0090	0	0		0	0		0	0		0	0	0	0	0	0	0	0		0	0	
1	0.0018	0	0		0	0		0	0		0	0	0	0	0	0	0	0		0	0	
2	0.0019	0	0		0	0		0	0		0	0	0	0	0	0	0	0		0	0	
3	0.0017	0	0		0	0		0	0		0	0	0	0	0	0	0	0		0	0	
1	negative control	0	0		0	0		0	0		0	0	0	0	0	0	0	0		0	0	
2	control	0	0		0	0		0	0		0	0	0	0	0	0	0	0		0	0	
3	control	0	0		0	0		0	0		0	0	0	0	0	0	0	0		1	0	
1	positive control	0	0		0	0		0	0		1	2	2	2	6	2,3	10	0		10	0	
2	control	1	0		1	1	3	2	0		3	5	2,3	5	2	3	9	1	3	10	0	
4	control	0	0		0	0		0	0		3	3	2	5	4	3	10	0		10	0	

Table 4 (continued): definitive oral toxicity test; mortality and behavioural abnormalities of the bees;

average ± standard deviation from three replicates per concentration / control.

avg. ± sd = average ± standard deviation; behav. abnorm. = behavioural abnormalities;

test article dosage avg. µg/bee	first 30 minutes				after 1 hour				after 4 hours				after 24 hours				after 48 hours			
	mortality		behav.abnor.		mortality		behav.abnor.		mortality		behav.abnor.		mortality		behav.abnor.		mortality		behav.abnor.	
	avg. ± sd		avg. ± sd		avg. ± sd		avg. ± sd		avg. ± sd		avg. ± sd		avg. ± sd		avg. ± sd		avg. ± sd		avg. ± sd	
	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%
0.169	0.0	0.0	0.0	0.0	0.0	0.0	23.3	15.3	0.0	0.0	56.7	20.8	40.0	20.0	6.7	5.8	80.0	26.5	0.0	0.0
0.083	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	6.7	11.5	3.3	5.8	20.0	10.0	0.0	0.0	23.3	5.8	0.0	0.0
0.0173	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	6.7	5.8	0.0	0.0	6.7	11.5	0.0	0.0	10.0	10.0	0.0	0.0
0.0083	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
0.0018	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
negative	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3.3	5.8	0.0	0.0
positive	3.3	5.8	0.0	0.0	6.7	11.5	0.0	11.5	40.0	17.3	40.0	17.3	96.7	5.8	3.3	5.8	####	0.0	0.0	0.0

Dit document is geen eigendom van het Ctgb en wordt beschikbaar gemaakt op grond van een wettelijke verplichting tot openbaarmaking. Op dit document kunnen rechten van derden rusten, waaronder intellectuele eigendomsrechten en/of auteursrechten. Voorts kan dit document onder een regeling ontrent gegevensbescherming vallen. Het gebruik van de inhoud hiervan deelt de rechten van de rechthebbende.

This document is not the property of the Ctgb and only provided based on mandatory freedom of information requirements. Consequently, any publication, distribution and/or publishing and any commercial exploitation and copy rights of third parties without the permission of the owner of this document may therefore be prohibited.

table 5: definitive oral intake test: weight of syringes at start of the experiment and after removing from the test cages; uptaken solution as calculated average (difference/bee)

test substance dosage nominal	test substance concentrations in food $\mu\text{g} / \text{mg}$	weight of syringes		difference mg	uptaken solution mg / bee	uptaken test substance	
		start mg	end mg			$\mu\text{g} / \text{bee}$	σ $\mu\text{g} / \text{bee}$
0.1 $\mu\text{g}/\text{bee}$	1 0.005	3517	3174	343	34	0.172	0.169
	2 0.005	3713	3356	357	36	0.179	
	3 0.005	3794	3478	316	32	0.158	
0.05 $\mu\text{g}/\text{bee}$	1 0.0025	3527	3209	318	32	0.080	0.083
	2 0.0025	3748	3393	355	36	0.089	
	3 0.0025	3553	3233	320	32	0.080	
0.01 $\mu\text{g} / \text{bee}$	1 0.0005	3423	3079	344	34	0.0172	0.0173
	2 0.0005	3544	3184	360	36	0.0180	
	3 0.0005	3583	3252	331	33	0.0166	
0.005 $\mu\text{g} / \text{bee}$	1 0.00025	3541	3248	293	29	0.0073	0.0083
	2 0.00025	3581	3236	345	35	0.0086	
	3 0.00025	3646	3288	358	36	0.0090	
0.001 $\mu\text{g} / \text{bee}$	1 0.00005	3716	3366	350	35	0.0018	0.0018
	2 0.00005	3596	3216	380	38	0.0019	
	3 0.00005	3566	3236	330	33	0.0017	
negative control	1 0	3587	3287	300	30	0.000	0.0
	2 0	3545	3238	307	31	0.000	
	3 0	3515	3183	332	33	0.000	
positive control Dimethoate 0.2 $\mu\text{g}/\text{bee}$	1 0.01	3620	3273	347	35	0.347	0.33
	2 0.01	3479	3148	331	33	0.331	
	3 0.01	3383	3065	318	32	0.318	