




TITLE PAGE

Residue Levels of Imidacloprid and Imidacloprid Metabolites in Nectar, Blossoms and Pollen of Summer Rape Cultivated on Soils with Different Imidacloprid Residue Levels and Effects of These Residues on Foraging Honeybees

Test Location: farmland "Laacher Hof" - 1999

**Report Amendment No. 1
13. April 2007**

AUTHOR



TESTING FACILITY

BAYER AG

**Crop Protection-Development
Institute For Environmental Biology**

D-51368 Leverkusen-Bayerwerk

GLP STUDY NUMBER

E 370 1548 - 8

REPORT NUMBER

SXR/Am 008

TOTAL NUMBER OF PAGES

2

STUDY COMPLETION DATE:

28 September 1999

Reason for this amendment:

Correction of a typing error on Page 9, section 3.7:

In the case of "Variant 1999", the test item was misreported due to a typing error. Summer rape seeds were treated with Poncho FS 500 instead of Gaucho WS 70.

Original version:

- Variant „1999“: untreated grass area since 1996. Drilled with Gaucho WS 70 treated summer rape seed on 12 May 1999 (33.5 g ai/ha).

Corrected version:


- Variant „1999“: untreated grass area since 1996. Drilled with Poncho FS 500 treated summer rape seed on 12 May 1999 (33.5 g ai/ha).

Impact on the Outcome of the Study:

None

DEPUTY OF STUDY DIRECTOR

Bayer CropScience AG:




Date: 2007-04-13

BCS-D-ETX

SPONSOR

BAYER CropScience AG:




Date: 2007-04-13,

Manager of the Testing Facility
(BCS-D-ETX)

and Representative of the Sponsor

GLP QUALITY ASSURANCE

BAYER CropScience AG:



Date: 2007-04-16

(BCS-D-GLP/QA)

TITLE PAGE

Residue Levels of Imidacloprid and Imidacloprid Metabolites in Nectar, Blossoms and Pollen of Summer Rape Cultivated on Soils with Different Imidacloprid Residue Levels and Effects of These Residues on Foraging Honeybees

Test Location: farmland "Laacher Hof" 1999

AUTHOR



TESTING FACILITY

BAYER AG

Crop Protection-Development
Institute For Environmental Biology
D-51368 Leverkusen-Bayerwerk

GLP STUDY NUMBER

E3701548 - 8

REPORT NUMBER

SXR/Am 008

TOTAL NUMBER OF PAGES

43

STUDY COMPLETION DATE:


28 September 1999



SXR/Am 008 / MO-99-015677

STATEMENT OF COMPLIANCE

This study was conducted in compliance with the Principles of Good Laboratory Practice (Chemicals Law (ChemG) of July 25, 1994, Annex 1 and OECD Principles of Good Laboratory Practice (GLP) of November 26, 1997 [C(97) 186/Final]).



Study Director

Title

28.9.99

Date

Responsible Analyst
Biological Samples

Title

28.9.99

Date

Responsible Analyst
Soil Samples

Title

28.9.99


Date

Dit document is geëigend
Op dit document is geen aanspraak
Publicatie, verspreiden of anderszins openbaar maken is
rechthebbende.

This document is not the property of the Ctgb and only provided based on mandatory freedom of information requirements.
The document may be subject to rights such as intellectual property and copy rights of third parties.
Consequently, any publication, distribution, reproduction and/or publishing and any commercial exploitation and use of this document or its contents without the permission of the owner of this document may therefore be prohibited and violate the rights of its owner.

af gemaakt op grond van een wettelijke verplichting tot openbaarmaking.
waaronder intellectuele eigendomsrechten en/of auteursrechten.
een regeling omtrent gegevensbescherming vallen.
exploitatie en gebruik van dit document of de inhoud hiervan zonder de toestemming van de rechthebbende.
e verboden zijn en een inbreuk opleveren van de rechten van deze rechthebbende.
exploitatie en gebruik van dit document of de inhoud hiervan zonder de toestemming van de rechthebbende.
exploitatie en gebruik van dit document of de inhoud hiervan zonder de toestemming van de rechthebbende.
exploitatie en gebruik van dit document of de inhoud hiervan zonder de toestemming van de rechthebbende.

CERTIFICATION OF AUTHENTICITY



Study Director

28.9.99

Title

Date

Responsible Analyst
Biological Samples

28.9.99

Title

Date

Responsible Analyst
Soil Samples

28.9.99

Title

Date

Head of Institute for
Environmental Biology

28.9.99

Title

Date

INQUIRIES

Inquiries should be directed to:



BAYER AG
Crop Protection - Development
Institute for Environmental Biology
D-51368 Leverkusen - Bayerwerk, FRG

Phone No. (02173) / 38 - 3888

Dit document is geen publicatie, verspreiding, of gebruik van dit document of de inhoud hiervan zonder de toestemming van de rechthebbende. Dit document is not the publication, distribution, reproduction or use of this document or its contents without the permission of the owner of this document. Furthermore, any publication, distribution, reproduction or use of this document or its contents may fall under a regulatory data protection regime. The document may be subject to rights such as intellectual property and copyright of third parties. This document is not the publication, distribution, reproduction or use of this document or its contents without the permission of the owner of this document. Furthermore, any publication, distribution, reproduction or use of this document or its contents may fall under a regulatory data protection regime. The document may be subject to rights such as intellectual property and copyright of third parties. This document is not the publication, distribution, reproduction or use of this document or its contents without the permission of the owner of this document. Furthermore, any publication, distribution, reproduction or use of this document or its contents may fall under a regulatory data protection regime. The document may be subject to rights such as intellectual property and copyright of third parties.

TABLE OF CONTENTS

	Page No.
TITLE PAGE.....	1
STATEMENT OF COMPLIANCE.....	2
CERTIFICATION OF AUTHENTICITY.....	3
TABLE OF CONTENTS.....	4
1.0 SUMMARY.....	5
2.0 INTRODUCTION.....	7
3.0 EXPERIMENTAL.....	7
3.1 Test Substance Used for Test Scenario 1999.....	7
3.2 Reference Substance.....	7
3.3 Execution of the Test.....	8
3.4 Origin of Honeybees and Preparation of Hive Colonies.....	8
3.5 Procedure of Seed Dressing.....	8
3.6 Location of the Trial Site and Description of the Study Plots.....	8
3.7 Treatment Design.....	9
3.8 Plot History and Cultivation of the Plots during the Study.....	10
3.9 Sampling Procedure.....	10
3.10 Sample Processing and Residue Analysis.....	11
3.11 Climatic Conditions During the Study.....	11
3.12 Observations on Honeybees.....	11
4.0 FILING.....	12
5.0 RESULTS AND DISCUSSION.....	13
5.1 Analytical Findings.....	13
5.2 Biological Observations on Foraging Honeybees.....	13
FIGURES.....	14
Figure 1: Arrangement of study plots on the study fields.....	14
TABLES.....	15
Table 1: Soil Residue Level of Imidacloprid at the Different Study Sites.....	15
Table 2: Biological Observations on the Hive Colonies Used for Sampling Summer Rape Nectar and Pollen.....	16
APPENDICES.....	20
APPENDIX I: Analytical Report for Soil Samples.....	20
APPENDIX II: Plot History and Cultivation of the Plots during the Study.....	27
APPENDIX III: Analytical Report for Biological Samples.....	31
Appendix IV: Copy of the GLP Certificate.....	41
Appendix V: Quality Assurance Statement.....	43

1.0 SUMMARY

Report: [REDACTED] (1999): Residue Levels of Imidacloprid and Imidacloprid Metabolites in Nectar, Blossoms and Pollen of Summer Rape Cultivated on Soils with Different Imidacloprid Residue Levels and Effects of These Residues on Foraging Honeybees. Test Location: farmland "Laacher Hof" - 1999
Bayer AG, unpublished report No: SXR/Am 008; 1999/09/28.
(appendix I and III report data from study MR471-99 and MR-518/99, respectively).

Guidelines: Internal Testing Method
Deviations: not applicable

GLP: yes (certified laboratory)

Material and methods: summer rape seed (variety "Lisonne") either dressed with 25 ml/kg Poncho FS 500 (a.i. content: 79.7 g/L beta-Cyfluthrin and 427.4 g/L imidacloprid; batch no. 6200/0055*A according to formulation no. 6200/0059, developmental no. 00195939) or imidacloprid-free were drilled on 12 May 99 in soils with different imidacloprid residue levels. Soil samples for an analytical determination of the imidacloprid residue level were taken immediately before drilling. Drilling rate was 3.25 kg/ha. During peak flowering of the summer rape (mid of July) small bee colonies (2,000 to 3,000 honeybees) were caged on these plots (appr. 50 m²) as a sampling device for summer rape nectar and pollen. In addition, some nectar and flowers were sampled by hand. The honeybees used as samplers were observed for signs of behavioral impacts. All samples and a small sample of honeybees were subjected to a residue analysis for imidacloprid and its relevant metabolites.

Dates of biological work: July 12 – 20, 1999
Dates of soil analysis: August 9 – 11, 1999
Dates of analysis of biological samples: August 30 – September 21, 1999.

Findings: Residues in soil, in summer rape plant matrices planted as succeeding crop and in honeybees used as sampling device.
(detects above the LOQ are highlighted):

Type of Sample	Residue Level [mg/kg] *		
	Imidacloprid	Olefin-NTN	Hydroxy-NTN
Control Plot (field number 711)			
Soil sample (0-30 cm)	n.d.	--	--
Leaves (produced latest)	n.d.	n.d.	n.d.
Flowers	n.d.	n.d.	n.d.
Nectar sampled from the flowers	n.d.	n.d.	n.d.
Pollen sampled from hives and bees	n.d.	n.d.	n.d.
Honeybees exposed to the summer rape	n.d.	n.d.	n.d.

* Limit of quantitation for soil samples: 0.006 mg/kg for imidacloprid; n.d. = below limit of detection (0.002 mg/kg)
Limit of quantitation for biological samples: 0.005 mg/kg for imidacloprid and hydroxy-imidacloprid, 0.01 mg/kg for olefin-imidacloprid. n.d. = below limit of detection (0.0015 and 0.003 mg/kg).

Type of Sample	Residue Level [mg/kg] *		
	Imidacloprid	Olefin-NTN	Hydroxy-NTN
Variant „1997“ (field number 710)			
Soil sample (0-30 cm)	0.016	--	--
Leaves (produced latest)	< LOQ	n.d.	n.d.
Flowers	n.d.	n.d.	n.d.
Nectar sampled from the flowers	n.d.	n.d.	n.d.
Pollen sampled from hives and bees	n.d.	n.d.	n.d.
Honeybees exposed to the summer rape	n.d.	n.d.	n.d.
Variant „1998“ (field number 702)			
Soil sample (0-30 cm)	0.013	--	--
Leaves (produced latest)	< LOQ	n.d.	n.d.
Flowers	n.d.	n.d.	n.d.
Nectar sampled from the flowers	n.d.	n.d.	n.d.
Pollen sampled from hives and bees	< LOQ	n.d.	n.d.
Honeybees exposed to the summer rape	n.d.	n.d.	n.d.
Variant „1998 (2x)“ (field number A XII)			
Soil sample (0-30 cm)	0.014	--	--
Leaves (produced latest)	< LOQ	n.d.	< LOQ
Flowers	n.d.	n.d.	n.d.
Nectar sampled from the flowers	n.d.	n.d.	n.d.
Pollen sampled from hives and bees	< LOQ	n.d.	n.d.
Honeybees exposed to the summer rape	n.d.	n.d.	n.d.
Variant „1999“ (field number 711)			
Soil sample (0-30 cm)	n.d.	--	--
Leaves (produced latest)	< LOQ	n.d.	< LOQ
Flowers	n.d.	n.d.	n.d.
Nectar sampled from the flowers	< LOQ	n.d.	n.d.
Pollen sampled from hives and bees	n.d.	n.d.	n.d.
Honeybees exposed to the summer rape	n.d.	n.d.	n.d.

* Limit of quantitation for soil samples: 0.006 mg/kg for imidacloprid; n.d. = below limit of detection (0.002 mg/kg)

Limit of quantitation for biological samples: 0.005 mg/kg for imidacloprid and hydroxy-imidacloprid, 0.01 mg/kg for olefin-imidacloprid. n.d. = below limit of detection (0.0015 and 0.003 mg/kg).

Observations: No treatment-related behavioral impacts (e.g. apathy, exaggerated motility, dis-coordinated movements) or suspicious mortality was observed on the honeybee colonies used for collecting summer rape nectar and pollen. The small colonies were remained till 3 September. The final check on this day did also reveal no abnormalities in either colony strength or brood status.

2.0 INTRODUCTION

According to EU directive 91/414/EEC the impacts of pesticides on honeybees have to be examined. Besides the intrinsic toxicity of a pesticide the concentration to which a honeybee may be exposed under field conditions is an integral component for the hazard assessment. The present study aims to examine the exposure in greater detail for a refined risk assessment.

The summer rape samples were analysed for residues of imidacloprid and its olefin- and hydroxy-metabolites. These metabolites were considered as relevant, since they have a chemical structure closely related to the parent molecule and were observed in plant metabolism studies in significant proportions (up to approx. 10 %).

3.0 EXPERIMENTAL

3.1 Test Substance Used for Test Variant „1999“

Test substance:	Poncho FS 500
Active ingredient(s):	(a) Beta-Cyfluthrin (FCR 4545) (b) Imidacloprid (NTN 33893)
Chemical name(s) of ai(s):	(a) Cyclopropanecarboxylic acid, 3-(2,2-dichloroethenyl)-2,2-dimethyl-, cyano (4-fluoro-3-phenoxyphenyl)methyl ester (b) 2-Imidazolidinimine, 1-[(6-chloro-3-pyridinyl)methyl]-N-nitro-
CAS number of ai(s):	(a) 68 359-37-5 (b) 138 261-41-3
Indikation:	seed dressing
Developmental/article number	0195939
Formulation/batch number:	6200/0055*A according to formulation no. 6200/0059
No. of certificate:	FAR-No. 642-00
AI content (acc. to analysis):	(a) 79.7 g/L (b) 427.4 g/L
Analytical method:	(a) GC, int. std. (b) HPLC, ext. std.
Date of analysis:	March 17, 1999
Expiry date:	March 17, 2000
Physical appearance:	darkblue suspension
Specific density:	1.155 g/ml
Storage conditions:	room temperature
Seed dressing rate(s) tested in the study:	25 ml/kg Poncho FS 500 (= nominal content: 2 g/kg beta-cyfluthrin & 10.6 g/kg imidacloprid; analytical findings, FAR 670-00: 1.8 g/kg beta-cyfluthrin & 10.3 g/kg imidacloprid).
Seed drilling rate tested in the study:	3.25 kg/ha (= 20,202 seed per four 240 m ² study plots) (summer rape variety: „Lisonne“; TGW: 3.86 g; standard fungicidal treatment: TMTD)
Safety precaution:	Routine hygienic precautions

3.2 Reference Substance

For this type of material and use pattern, a reference compound is not specified.

3.3 Execution of the Test

The sampled study plots were drilled on 12 May 1999. Sampling of nectar, pollen, flowers and honeybees and the behavioral observations were performed between 6 and 20 July 1999.

Sponsor:

BAYER AG
GB Plant Protection
Marketing - Seed Treatment
D-40789 Monheim

Study Director:

Cultivar Manager:

Trials Officer:

Responsible Analyst (soil)

Responsible Analyst (biological samples):

Study Technicians:

Quality Assurance:

Laboratory Study Number:

SXR/Am 008

3.4 Origin of Honeybees and Preparation of Hive Colonies

Honeybee colonies used for pollen and nectar sampling were supplied by a German beekeeper [REDACTED]. Preparation of the hive colonies used for the test started on 22 May 1999. From large commercially managed beehives, 20 combs were removed and combined. After the brood of these combs had ecdysed (5 June 1999), the combs were removed and allocated to 10 three-comb-colonies (two combs from the combined hive and a new comb matrix) with about 2,000 – 3,000 workerbees. Two days before colony installment on the test plots, the two original combs were replaced by new comb matrices and the colonies received 0.5 litre ready-to-use syrup (1:1). Hive installment was on 10 July 1999. The small bee colonies were retained on the study plots till 20 July 1999.

3.5 Procedure of Seed Dressing

The summer rape seeds (variety: „Lisonne“) used for test “variant 99” were dressed by a commercial seed dressing company (SUET Saat- und Erntetechnik GmbH, D-37257 Eschwege) and delivered to Bayer on 1 April 1999. Besides the insecticidal treatment, the seed were treated with a standard fungicide (TMTD). This fungicidal treatment was also applied to all imidacloprid-free seeds which were drilled on study plots of test “variants 97, 98, 98 (2x)” and the control.

3.6 Location of the Trial Site and Description of the Study Plots

The trial site was at Bayer AG’s experimental farmland "Laacher Hof", approximately 3 km south of Monheim (Germany, NRW 41m above sea level). The precise plot location was as follows:

- Control plot: field number 711
- Variant „1997“: field number 710
- Variant „1998“: field number 702

- Variant „1998 (2x)“: field number A XII
- Variant „1999“: field number 711

The soil characteristics of the study plots were determined for another study for a field close to the study fields (OE No. 2565, sampling date: 8 December 1998). The soil at this site was classified as a "silty sand" with particle size fractions of 78.5% sand, 19.2% silt and 2.3% clay. The pH value (KCL) at the study site was determined to be 6.08. Soil organic carbon was 0.75% by weight. The water holding capacity was 39.08 g water per 100 g dry soil.

3.7 Treatment Design

After the previous crop had been destroyed (4 l/ha Glyphos and subsequent ploughing), all study plots were drilled with 3.25 kg/ha summer rape seed on 12 May 1999. For each test variant and for the control, plots of 8 x 30 m were drilled with either imidacloprid-free or Poncho FS 500 dressed summer rape seed (variety: „Lisonne“). Drilling distance was 21.6 cm between rows and 5.5 cm in-row. Prior to sowing the proper functioning of the seeding equipment was tested and adjusted to the target conditions (e.g. seed density, in-row distances,). The test plots were adjacent to similar test plots which were cultivated with either maize or sunflower plants.

With regard to imidacloprid, study plots received since 1996 the following treatments¹:

- Control plot: untreated grass area since 1996. Drilled with imidacloprid-free summer rape seed on 12 May 1999
- Variant „1997“: cropped in fall 1997 with Gaucho treated winter wheat (77 g ai/ha), sprayed on 24 April 1999 with 71.5 g/ha Gaucho WS 70 (= 50 g ai/ha imidacloprid; batch no. 233 614 749, 72.5% imidacloprid according to FAR no. 559-01). Drilled with imidacloprid-free summer rape seed on 12 May 1999.
- Variant „1998“: cropped in spring 1996 with Gaucho treated sugar beet (111 g ai/ha), followed in fall 1998 by Gaucho treated winter barley (49 g ai/ha). Drilled with imidacloprid-free summer rape seed on 12 May 1999.
- Variant „1998 (2x)“: cropped in spring 1998 with Gaucho treated sugar beet (105 g ai/ha) followed by Gaucho-treated winter wheat (76 g ai/ha). Drilled with imidacloprid-free summer rape seed on 12 May 1999.
- Variant „1999“: untreated grass area since 1996. Drilled with Gaucho WS 70 treated summer rape seed on 12 May 1999 (33.5 g ai/ha).

On the day of drilling, soil samples were taken to analytically verify the residue level of the study plots. From each study field 20 soil cores of 5 cm diameter and a depth of 30 cm were sampled. Sampling points were distributed along the two diagonals of each study field with equal distances between the points, i.e. 10 samples per diagonal.

Depending on the plot arrangement, the total size of the sampled area was:

- Control plot/Variant „1999“: 30 x 50 m.
- Variant „1997“: 24 x 30 m.
- Variant „1998“: 24 x 30 m.
- Variant „1998 (2x)“: 8 x 90 m.

Immediately after sampling, soil samples were divided into two subsamples, one subsample contained the 0-20 cm top soil layer and the other subsample the 20-30 cm soil fraction. After dividing, all subsamples were stored at -20°C until residue analysis.

¹ Crop management before 1999 was not conducted and recorded under GLP regulations.

Residue levels of the different subsamples are reported in the pertinent analytical report (appendix I).

Shortly before full rape blossom, tunnel cages of 10 x 5 m and 3 m height were installed on each study plot. The tunnel cages consisted of an aluminium frame covered by plastic gauze material (2 x 2 mm mesh size). For operational purposes, a walkway was created by removing all plants along a 50 cm wide transect from the tunnel entrance to the opposite end.

3.8 *Plot History and Cultivation of the Plots during the Study*

Plot history and 1999 treatments of the study plots are reported in detail in appendix II.

3.9 *Sampling Procedure*

Installment of bee colonies

One bee hive with 3 combs (about 2,000 – 3,000 honeybees) was placed left from the tunnel entrance on each study plot on 10 July 1999 (hive colonies no. 45, 40, 56, 58, 44). Colonies remained in the tunnels till 20 July 1999.

Sampling of Foraging Honeybees

On days 7 and 9 following colony installment (= 17 and 19 July 1999) approximately 100 honeybees were sampled with glass tubes while foraging on the summer rape within the tunnel. Sampled honeybees were killed by placing the sampling glass tubes into a container with dry ice. At the end of each sampling day at the latest, collected honeybees were transferred into a refrigerator (-20°C) where they were retained until residue analysis (see 3.10). This sampling was primarily done to get summer rape nectar from the honeybulbs and summer rape pollen from the pollen pockets. In terms of time and sample quantity, however, other sampling methods (see below) proved to be more efficient. For this reason, no further processing of honeybees was done as initially intended but the honeybees kept in reserve in case that other questions may arise during or after the analytical work. A small number of honeybees (approximately 1 g) were analyzed for residues to examine whether or not any residues in nectar and/or pollen would also be detected in this matrix.

Sampling of Summer Rape Nectar

Between 6 and 11 July 1999 (= 54 to 59 days after seed drilling) nectar from rape plants was sampled directly from the flowers with micro-capillaries. In preparation to this sampling procedure, sampling plots were confined with small gauze-covered tents (2 x 2 m) shortly before flowering to exclude other nectar-feeding arthropods and thus to avoid exploitation of this nectar source. Nectar samples were emptied from the microcapillaries into small plastic tubes (1.5 mL Eppendorf vials) till a minimum sample volume of 0.5 mL was harvested. During sampling, the plastic tubes with the nectar samples were stored on dry ice in the field. At the end of each sampling day at the latest, the nectar samples were transferred into a refrigerator (-20°C) where they were retained until residue analysis (see 3.10).

Sampling of Summer Rape Pollen from Honeybees and from the Beehives

On the day of hive removal (= 20 July 1999) combs were taken off the beehives to cut out pollen stores. Additional all pollen pockets were sampled from the 100 honeybees collected while foraging on the summer rape within the tunnel plots. After sampling

(either from combs or from honeybees), the pollen samples were stored on dry ice. At the end of each sampling event at the latest, the pollen samples were transferred into a refrigerator (-20°C) where they were retained until residue analysis (see 3.10).

Sampling of Summer Rape Flowers

About 20 g of each, male and female flowers were sampled from the summer rape plants during the peak flowering period. After sampling, the flowers were stored on dry ice in the field. At the end of each sampling day at the latest, the flower samples were transferred into a refrigerator (-20°C) where they were retained until residue analysis (see 3.10).

Sampling of Summer Rape Leaves

Sunflower leaves were collected on 13 July 1999. After sampling, the leaves were stored on dry ice in the field. At the end of the sampling day, the leaf samples were transferred into a refrigerator (-20°C) where they were retained until residue analysis (see 3.10).

3.10 Sample Processing and Residue Analysis

Sample processing and analytical methods are described in detail in appendix I (soil samples) and appendix III (biological samples).

3.11 Climatic Conditions During the Study

During cultivation of the study plots, temperature and precipitation events were continuously recorded by weather stations located adjacent to the study sites (within a 3 km distance). The following records were made during this time period:

Month	Precipitation [mm]	Min. air temperature 2m [°C]	Max. air temperature 2m [°C]	Soil temperature 0 cm [°C]	Energy input [kJ/cm ²]
April	70.9	1.2 – 12.4	7.4 – 22.7	3.8 – 13.4	39.9
May	49.6	6.3 – 17.9	14.8 – 30.4	12.3 – 25.9	54.5
June	71.8	9.2 – 17.2	15.6 – 30.3	13.0 – 26.6	52.4
July	41.1	13.1 – 20.7	18.8 – 33.0	16.2 – 30.0	59.1
August	77.8	8.6 – 19.9	18.7 – 32.4	14.1 – 28.1	46.0

While hive colonies were confined within the tunnels, climatic records were made during each evaluation. The following conditions were recorded (checks between high noon till 4 p.m):

Day after first exposure	Precipitation [mm]	Air temperature [°C]	Soil temperature [°C]	Cloudness [%]	Wind Relations*
2	2	24	26	90	-
3	2	26	30	80	-
4	5	20	22	80	++
5	1	20	22	95	+
6	0	24	26	50	++
9	0	32	34	0	-

- calm, + slight, ++ moderate, +++ strong

3.12 Observations on Honeybees

All behavioral anomalies of the honeybees were recorded together with the date of observations. In particular, the following behavioural components were noted:

- Foraging intensity:** Six times (days 2, 3, 4, 5, 6 and 9 after 1st exposure) the number of bees foraging during 1 minute within a randomly selected 1 m² area of the summer rape plot was recorded.
- Behavioral Anomalies:** Whenever observed, the following behavioral anomalies were recorded with the date and daytime of observation:
- exaggerated motility
 - discoordinated movements (trembling, shaking)
 - apathy.
- Mortality:** In front of the hive colonies, linen sheets of 60 x 50 cm were placed on the ground to trap the dead bees which were removed from the beehives during the time while colonies were confined within the tunnel cages. In addition, the number of dead honeybees around the tunnel margin were counted as an indicator whether a higher number of bees tried to leave the tunnel or failed to return to the hive. Also, any conspicuous numbers of dead bees in the study plots was recorded but no formal counts were made.
- Colony strength and brood status:** Colonies were checked on 20 July for colony strength and brood status.

4.0 FILING

All raw data, the study protocol and the original of the report are filed in the Central GLP archive of PF/E, Crop Protection Center 40789 Monheim, FRG. Reserve samples of the test substance are stored in the pertinent archive of that test facility which provided or certified the test substance.

Dit document is geen eigendom van het Ctgb en wordt beschikbaar gemaakt op grond van de Wet van 1992 betreffende de openbaarheid van bestuur.
Op dit document kunnen rechten van auteurs, uitgeverijen, waardenrechten, waaronder intellectuele eigendomsrechten, voortvloeien.
Voorts kan dit document in het openbaar gebruik van dit document een inbreuk opleveren op de rechten van de rechthebbende van dit document kan derhalve verboden zijn.
This document is not the property of the Ctgb and only provided based on mandatory freedom of information provisions.
The document may be subject to rights such as intellectual property and copy rights of third parties.
Furthermore, this document may fall under a regulatory data protection regime.
Consequently, any publication, distribution, reproduction and/or publishing and any form of commercial exploitation and use of this document or its contents without the permission of the owner of this document may therefore be prohibited and violate the rights of its owner.

5.0 RESULTS AND DISCUSSION

5.1 Analytical Findings

Soil Samples

Analytical findings are summarized in table 1 and given in detail in the analytical report (appendix I). In the control plots, no residues at or above the limit of quantitation was detected. In the treated plots, residue levels were well in the range as expected from plot and/or treatment history. Within the 0-30 cm soil layer, imidacloprid concentrations of 15.7, 14.3 and 12.7 µg/kg were determined for the test variants „1997“, „1998 (2x)“, and „1998“, respectively (Tab. 1). No residues were detected in the 0-30 cm soil samples of the „1999/control“ field.

Biological Samples

Residue levels of imidacloprid and of its toxicologically relevant metabolites (olefin- and hydroxy-imidacloprid) were below the limit of quantitation (= 0.005 and 0.01 mg/kg, respectively) in all bee-relevant summer rape parts (nectar, pollen), in the summer rape flowers and the sampling honeybees as well. Residues slightly above the limit of detection were found in the nectar sample of test variant „1999“, i.e. from the seed-treated plants, and in the pollen samples of test variants „1998“ and „1998 (2x)“. The youngest rape leaves of all treatment sites had residue levels of imidacloprid above the limit of detection but below the limit of quantitation. No residues were found in the control leaves. In the test variants „1998 (2x)“ and „1999“, traces of the hydroxy-metabolite (less than 5 µg/kg) were detected in the young leaves.

5.2 Biological Observations on Foraging Honeybees

All hive colonies decreased in weight during their confinement on the study plots (Tab. 2). The weight decrease was 2.9, 4.2, 2.9, 3.2 and 3.0% of the initial weight for the control, test variants „1997“, „1998“, „1998 (2x)“ and „1999“, respectively. These data show that the weight decrease was comparable between study plots and no treatment-related effect is indicated.

Foraging activity (recorded during 6 occasions between 12 and 19 July 1999) was 122, 82, 97, 99, and 134 honeybees, respectively, per 2 m² summer rape within a 1 minute observation period (Tab. 2). The differences in foraging activity are most likely attributed to the differences in the number of actually flowering summer rape plants within the different plots. Most summer rape plants were emerged and in flower on the control and test variant „1999“ which had the highest foraging activity as well.

Therefore, it can be concluded that there was no treatment-related effect on foraging activity.

During the foraging period, no behavioral impacts (e.g. apathy, exaggerated motility, disordinated movements) were observed.

The treatment-specific mortality figures were 11, 11, 9, 9 and 10 honeybees for the control and the test variants „1997“, „1998“, „1998 (2x)“ and „1999“, respectively (Tab. 2). No treatment-related impact on mortality can be concluded from these figures.

At study termination (20 July 1999), all colonies were finally checked for colony strength and brood status. There were no treatment-related differences in population density, food stores or brood status between the control and the treatment hives.

In summary, it can be concluded that honeybees were not adversely affected by any of the examined exposure scenarios.

FIGURES




Figure 1: Arrangement of the study plots on study fields.

Each study plot had a size of 8 x 30 m with distances of 21.6 cm between rows and 5.5 cm in-row.

Dit document is geen eigendom van het Ctgb en wordt beschikbaar gemaakt op grond van een wettelijke verplichting tot openbaarmaking.
 Op dit document kunnen rechten van derden rusten, waaronder intellectuele eigendomsrechten en/of auteursrechten.
 Voorts kan dit document onder een regeling omtrent gegevensbescherming vallen.
 Dit document is niet de eigendom van het Ctgb en wordt beschikbaar gemaakt op grond van een wettelijke verplichting tot openbaarmaking.
 The document may be subject to rights such as intellectual property and copy rights.
 Furthermore, this document may fall under a regulatory data protection regime.
 Consequently, any publication, distribution, reproduction and/or publishing and any commercial exploitation and use of this document or its contents without the permission of the owner of this document may therefore be prohibited and violate the rights of its owner.

TABLES

Table 1: Soil Residue Level of Imidacloprid at the Different Study Sites.

The details of the analytical work are given in appendix I. Residue data refer to the level immediately before seed drilling on 12 May 1999. Plot history was as follows:

- Control plot: last imidacloprid treatment: before 1996
- Variant „1997“: last imidacloprid treatment: 23 April 1999 (50 g ai/ha)
- Variant „1998“: last imidacloprid treatment: 24 Sept. 1998 (49 g ai/ha)
- Variant „1998 (2x)“: last imidacloprid treatment: 13 Oct. 1998 (76 g ai/ha)
- Variant „1999“: last imidacloprid treatment: before 1996, drilled on 12 May 1999 with Poncho-treated summer rape seed (33.5 g ai/ha).

Sample No.	Sample description	Soil Layer	Imidacloprid Residue Level [µg/kg]
1	Control Plot (field number 711)	0-30 cm	n.d.
2	Variant „1997“ (field number 710)	0-30 cm	15.7
3	Variant „1998“ (field number 702)	0-30 cm	12.7
4	Variant „1998 (2x)“ (field number A XII)	0-30 cm	14.3
5	Variant „1999“ (field number 711)	0-30 cm	n.d.

LOQ (Limit of quantification): 0.006 mg/kg.

n.d.: Residue levels below the limit of detection: 0.002 mg/kg.

Dit document is geen eigendom van het Ctgb en wordt beschikbaar gemaakt op grond van een wettelijke verplichting tot openbaarmaking. Op dit document kunnen rechten van derden rusten, waaronder intellectuele eigendomsrechten en/of auteursrechten. Voorts kan uit dit document onder een regeling omtrent gegevensbescherming vallen. Het gebruik van de inhoud hiervan zonder toestemming van de rechthebbende van dit document kan daartoe leiden tot een inbreuk op de rechten van derden.

This document is not the property of the Ctgb and only provided based on mandatory freedom of information requirements. The document may be subject to rights, such as intellectual property and copy rights of third parties. Consequently, any publication, distribution, reproduction and/or publishing and any commercial exploitation and use of this document or its contents without the permission of the owner of this document may therefore be prohibited and violate the rights of its owner.

Table 2: Biological Observations on the Hive Colonies Used for Sampling Summer Rape Nectar and Pollen.

Hive installment was on 10 July 1999. Plot history was as follows:

- Control plot: last imidacloprid treatment: before 1996
- Variant „1997“: last imidacloprid treatment: 23 April 1999 (50 g ai/ha)
- Variant „1998“: last imidacloprid treatment: 24 Sept. 1998 (49 g ai/ha)
- Variant „1998 (2x)“: last imidacloprid treatment: 13 Oct. 1998 (76 g ai/ha)
- Variant „1999“: last imidacloprid treatment: before 1996; drilled on 12 May 1999 with Poncho-treated summer rape seed (33.5 g ai/ha).

Weight Development

Treatment	Hive nucleus no.	Hive weight at study initiation [g]	Hive weight at study termination [g]	Weight changes [%]
Control	45	6785	6585	-2.9
Variant 97	40	6755	6470	-4.2
Variant 98	56	6655	6460	-2.9
Variant 98 (2x)	58	6560	6350	-3.2
Variant 99	44	6750	6545	-3.0

Foraging Activity

Treatment	Days after first exposure	No. honeybees foraging during 1 min within 1 m ²		Total no. foraging honeybees per plot
		Left of walkway	Right of walkway	
Control (= hive nucleus no. 45)	2	1	0	
	3	23	21	
	4	5	3	
	5	12	6	
	6	18	17	
	9	8	8	122
Variant „1997“ (= hive nucleus no. 40)	2	0	1	
	3	17	18	
	4	2	2	
	5	8	6	
	6	12	10	
	9	3	3	82
Variant „1998“ (= hive nucleus no. 56)	2	2	0	
	3	15	22	
	4	4	5	
	5	6	10	
	6	11	13	
	9	5	4	97

Table 2: cont'd.

Hive installment was on 10 July 1999. Plot history was as follows:

- Control plot: last imidacloprid treatment: before 1996
- Variant „1997“: last imidacloprid treatment: 23 April 1999 (50 g ai/ha)
- Variant „1998“: last imidacloprid treatment: 24 Sept. 1998 (49 g ai/ha)
- Variant „1998 (2x)“: last imidacloprid treatment: 13 Oct. 1998 (76 g ai/ha)
- Variant „1999“: last imidacloprid treatment: before 1996; drilled on 12 May 1999 with Poncho-treated summer rape seed (33.5 g ai/ha).

Treatment	Days after first exposure	No. honeybees foraging during 1 min within 1 m ²		Total no. foraging honeybees per plot
		Left of walkway	Right of walkway	
Variant „1998 (2x)“	2	0	0	
(= hive nucleus no. 58)	3	17	24	
	4	6	8	
	5	4	9	
	6	10	15	
	9	3	3	90
Variant „1999“	2	0	1	
(= hive nucleus no. 44)	3	20	25	
	4	6	7	
	5	9	8	
	6	20	20	
	9	10	8	134

Mortality

Treatment	Days after first exposure	No. of dead honeybees		No. of dead honeybees Total
		In front of beehive	At the tent margin	
Control	2	0	0	
	3	0	4	
(= hive nucleus no. 45)	4	0	3	
	5	0	2	
	6	0	1	
	9	0	1	11
Variant „1997“	2	0	0	
(= hive nucleus no. 40)	3	2	1	
	4	0	2	
	5	0	3	
	6	0	1	
	9	0	2	11
Variant „1998“	2	0	0	
(= hive nucleus no. 56)	3	1	0	
	4	0	2	
	5	0	2	
	6	0	2	
	9	0	2	9

Table 2: cont'd.

Hive installment was on 10 July 1999. Plot history was as follows:

- Control plot: last imidacloprid treatment: before 1996
- Variant „1997“: last imidacloprid treatment: 23 April 1999 (50 g ai/ha)
- Variant „1998“: last imidacloprid treatment: 24 Sept. 1998 (49 g ai/ha)
- Variant „1998 (2x)“: last imidacloprid treatment: 13 Oct. 1998 (76 g ai/ha)
- Variant „1999“: last imidacloprid treatment: before 1996; drilled on 12 May 1999 with Poncho-treated summer rape seed (33.5 g ai/ha).

Mortality

Treatment	Days after first exposure	No. of dead honeybees		No. of dead honeybees Total
		In front of beehive	At the tent margin	
Variant „1998 (2x)“ (= hive nucleus no. 58)	2	0	0	0
	3	0	1	1
	4	0	2	2
	5	0	2	2
	6	1	2	3
	9	0	1	1
Variant „1999“ (= hive nucleus no. 44)	2	0	0	0
	3	1	1	2
	4	0	2	2
	5	0	1	1
	6	0	2	2
	9	0	3	3
				10

Population Strength and Brood Status (20 July 1999)

Treatment	Occupation degree of central comb by honeybees	Brood development	Occupation of side combs by honeybees, food stores or brood
Control (= hive nucleus no. 45)	+++	++	-
Variant 97 (= hive nucleus no. 40)	++	++	-
Variant 98 (= hive nucleus no. 56)	+++	+++	-
Variant 98 (2x) (= hive nucleus no. 58)	+++	+++	+
Variant 99 (= hive nucleus no. 44)	+++	+++	-

- = no honeybees/food stores/brood cells on combs

+ = singular honeybees/food stores/brood cells on combs

++ = partly occupied combs (honeybees/food stores/brood cells)

+++ = well occupied combs (honeybees), good brood amount, high food stores.

Table 3: Plant Residue Level of Imidacloprid and Toxicologically Relevant Metabolites at the Different Study Sites.

The details of the analytical work are given in appendix III. Plot history was as follows:

- Control plot: last imidacloprid treatment: before 1996
- Variant „1997“: last imidacloprid treatment: 23 April 1999 (50 g ai/ha)
- Variant „1998“: last imidacloprid treatment: 24 Sept. 1998 (49 g ai/ha)
- Variant „1998 (2x)“: last imidacloprid treatment: 13 Oct. 1998 (76 g ai/ha)
- Variant „1999“: last imidacloprid treatment: before 1996; drilled on 12 May 1999 with Poncho-treated summer rape seed (3305 g ai/ha).

Type of Sample	Residue Level [mg/kg] *		
	Imidacloprid	Olefin-NTN	Hydroxy-NTN
Control Plot (field number 711)			
Leaves (produced latest)	n.d.	n.d.	n.d.
Flowers	n.d.	n.d.	n.d.
Nectar sampled from the flowers	n.d.	n.d.	n.d.
Pollen sampled from hives and bees	n.d.	n.d.	n.d.
Honeybees exposed to summer rape	n.d.	n.d.	n.d.
Variant „1997“ (field number 710)			
Leaves (produced latest)	< LOQ	n.d.	n.d.
Flowers	n.d.	n.d.	n.d.
Nectar sampled from the flowers	n.d.	n.d.	n.d.
Pollen sampled from hives and bees	n.d.	n.d.	n.d.
Honeybees exposed to summer rape	n.d.	n.d.	n.d.
Variant „1998“ (field number 702)			
Leaves (produced latest)	< LOQ	n.d.	n.d.
Flowers	n.d.	n.d.	n.d.
Nectar sampled from the flowers	n.d.	n.d.	n.d.
Pollen sampled from hives and bees	< LOQ	n.d.	n.d.
Honeybees exposed to summer rape	n.d.	n.d.	n.d.
Variant „1998 (2x)“ (field number A XII)			
Leaves (produced latest)	< LOQ	n.d.	< LOQ
Flowers	n.d.	n.d.	n.d.
Nectar sampled from the flowers	n.d.	n.d.	n.d.
Pollen sampled from hives and bees	< LOQ	n.d.	n.d.
Honeybees exposed to summer rape	n.d.	n.d.	n.d.
Variant „1999“ (south of field number 711)			
Leaves (produced latest)	< LOQ	n.d.	< LOQ
Flowers	n.d.	n.d.	n.d.
Nectar sampled from the flowers	< LOQ	n.d.	n.d.
Pollen sampled from hives and bees	n.d.	n.d.	n.d.
Honeybees exposed to summer rape	n.d.	n.d.	n.d.

* Limit of quantitation: 0.005 mg/kg (imidacloprid & hydroxy-metabolite), 0.01 mg/kg (olefin-metabolite); n.d. = below limit of detection (0.0015 mg/kg and 0.003 mg/kg, respectively)

APPENDICES

APPENDIX I: Analytical Report for Soil Samples.

Bayer AG, Crop Protection Business Group
Crop Protection - Development
Institute for Metabolism Research and Residue Analysis
51368 Leverkusen, Germany

August 31, 1999

MR-471/99

Page 20 of 7

Title**Analysis of Soil Samples from**

E 370 1548 - 8

E 370 1549 - 9

E 370 1550 - 0

E 370 1551 - 2

E 370 1552 - 3

E 370 1553 - 4

for Residues of Imidacloprid

Responsible Scientist

[REDACTED]
Bayer AG, Crop Protection Business Group
Crop Protection - Development
Institute for Metabolism Research and Residue Analysis (PF-E/MR)
51368 Leverkusen, Germany

Experimental Starting Date

August 09, 1999

Experimental Completion Date

August 11, 1999

Study Numbers

E 370 1548 - 8

E 370 1549 - 9

E 370 1550 - 0

E 370 1551 - 2

E 370 1552 - 3

E 370 1553 - 4

TABLE OF CONTENTS

	PAGE
TITLE	20
TABLE OF CONTENTS.....	21
1 INTRODUCTION.....	22
2 REFERENCE SUBSTANCE.....	22
3 PERFORMANCE	22
3.1 Extraction.....	22
3.2 High Performance Liquid Chromatographic Measurement.....	23
3.3 Method of Confirmation.....	23
4 RESULTS.....	24
5 REFERENCES.....	26

Dit document is geen eigendom van het Ctgb en wordt beschikbaar gemaakt op grond van een wettelijke verplichting tot openbaarmaking. Op dit document kunnen rechten van derden rusten, waaronder intellectuele eigendomsrechten en/of auteursrechten. Voorts kan dit document onder een regeling omtrent gegevensbescherming vallen. Het is niet toegestaan de inhoud hiervan zonder de toestemming van de rechthebbende van dit document te kopiëren, verspreiden of openbaar te maken. Het is niet toegestaan de rechten van de rechthebbende van dit document te openbaar te maken.

This document is not the property of the Ctgb and only provided based on mandatory freedom of information requirements. The document may be subject to rights such as intellectual property and copy rights of third parties. Consequently, any publication, distribution and/or publishing and any commercial exploitation and use of this document or its contents without the permission of the owner of this document may therefore be prohibited and violate the rights of its owner.


1 INTRODUCTION

Soil samples of the German trial stations "Höfchen" and "Laacher Hof" were analyzed for residues of Imidacloprid. The results are tabulated in Table 2 and 3. Extraction of soil samples and determination of Imidacloprid by HPLC-UV were performed according to method 00267 (MR-53/92) [3]. The limit of quantification (LOQ) was 6 µg/kg. The limit of detection (LOD) was 2 µg/kg.

2 REFERENCE SUBSTANCE

The following substance will be used as reference substance in recovery experiments and for preparation of external standard solutions.

Imidacloprid



Empirical formula:	$C_9H_{10}ClN_5O_2$
Molecular weight:	255.7 g/mol
Reference Substance No:	M00680
Purity:	99.4 % (HPLC), identity ensured by MS
Expiry date:	March 2000

3 PERFORMANCE

3.1 Extraction

Soil samples are extracted in a Soxtec extraction device with boiling methanol. The oil-bath temperature is set at 200 °C.

Soil samples of 25 g are weighed into an extraction thimble and covered with a defatted cotton wool plug. 40 mL of methanol and some boiling chips are placed into aluminum cups. Thimbles and cups are inserted in the Soxtec extraction device.

The extraction time takes one hour. Afterwards the thimbles are placed in rinse position for 30 minutes until the extraction is terminated.

The residue is flushed quantitatively into a 50 mL centrifuge tube by two times rinsing the aluminium cups with about 5 mL of ethanol. The extract is evaporated to dryness in a Turbo-Vap evaporator at 50 °C and reconstituted in 2 mL of acetonitrile/water 50/50 (v/v).

3.2 High Performance Liquid Chromatographic Measurement

Liquid chromatograph: Hewlett Packard 1090

Column: LiChrospher 60 RP-Select B (5 μ m) 125 x 4 mm

Solvent A: Water + 1g Sodium-dihydrogenphosphate-2-hydrate per L

Solvent B: Acetonitrile

Oven temperature: 40 °C

Inject. volume: 25 μ L

Flow rate: 1.5 mL/min

Detector wavelength: 270 nm

Table 1: Gradient for the HPLC-UV measurement

Time	0 min	10 % B
10 min.		25 % B
13 min		90 % B
18 min		90 % B
20 min		10 % B
30 min		10 % B

Retention time of Imidacloprid: approx. 6.4 min

3.3 Method of Confirmation

Within each series of analyses the identity of Imidacloprid was determined by LC/MS/MS according to method 00537 (MR-551/98) [4]. Therefore, one standard sample (recovery experiment), one control sample and one sample from the trials were analysed for the characteristic mass-to-charge ratio of Imidacloprid.

4 Results

Table 2: Concentrations of Imidacloprid for trial station "Höfchen"
(E3701551-2, E3701552-3 and E3701553-4)

Sample No.	Sample description	Soil layer	Imidacloprid [µg/kg]
No.1	Control sample (identical with test sample 1999)	0-20 cm	< LOQ
No.2	Control sample (identical with test sample 1999)	0-30 cm	n.d.
No.3	Test sample 1998	0-20 cm	8.7
No.4	Test sample 1998	0-30 cm	< LOQ
No.5	Test sample 1997	0-20 cm	24.5
No.6	Test sample 1997	0-30 cm	17.8

< LOQ: Concentrations of Imidacloprid below the limit of quantification of the analytical method of 6 µg/kg.

n.d.: Concentrations of Imidacloprid below the limit of detection of the analytical method of 2 µg/kg.

Dit document is geen eigendom van het Ctgb en wordt beschermd op grond van een wettelijke verplichting tot openbaarmaking.
 Op dit document kan de rechten van derden rusten, waaronder intellectuele eigendomsrechten en/of auteursrechten.
 Publicatie, verspreiding, vermenigvuldigen, verspreiden of anderszins openbaar maken van dit document of de inhoud hiervan onder de toestemming van de rechthebbende.
 This document is not the property of the Ctgb and only provided based on mandatory freedom of information requirements.
 The document may be subject to rights of intellectual property and copy rights of third parties.
 Consequently, any publication, distribution, reproduction and/or publishing and any commercial exploitation and use of this document or its contents without the permission of the owner of this document may therefore be prohibited and violate the rights of its owner.

Table 3: Concentrations of Imidacloprid for trial station “Laacher Hof” (E3701548-8, E3701549-9 and E3701550-1)

Sample No.	Sample description	Soil layer	Imidacloprid [µg/kg]
No.1	Control sample (identical with test sample 1999)	0-20 cm	< LOQ
No.2	Control sample (identical with test sample 1999)	0-30 cm	n.d.
No.3	Test sample 1998	0-20 cm	15.3
No.4	Test sample 1998	0-30 cm	12.7
No.5	Test sample 1998 (replicate)	0-20 cm	16.1
No.6	Test sample 1998 (replicate)	0-30 cm	14.3
No.7	Test sample 1997	0-20 cm	17.3
No.8	Test sample 1997	0-30 cm	15.7

< LOQ: Concentrations of Imidacloprid below the limit of quantification of the analytical method of 6 µg/kg.

n.d.: Concentrations of Imidacloprid below the limit of detection of the analytical method of 2 µg/kg.

Dit document is een eigendom van het Ctgb en wordt beschikbaar gemaakt op grond van een wettelijke verplichting tot openbaarmaking. Op dit document kunnen rechten van derden rusten, waaronder intellectuele eigendomsrechten en/of auteursrechten. Publicatie, verspreiding, vermenigvuldiging, commercieële exploitatie en gebruik van dit document of de inhoud hiervan zonder de toestemming van de rechthebbende zijn een inbreuk opleveren van de rechten van de rechthebbende.

This document is not the property of the Ctgb and only provided based on mandatory freedom of information requirements. The document may be subject to rights such as intellectual property and copy rights of third parties. Furthermore, reproduction and use of this document or its contents without the permission of the owner of this document may therefore be prohibited and violate the rights of its owner.

Table 4: Recovery Rates of Imidacloprid

Fortification [µg/kg]	Soil	Soil layer	Imidacloprid [%]
6.02	Höfchen	0-20 cm	94.0
6.02	Laacher Hof	0-20 cm	95.7
60.2	Höfchen	0-20cm	92.2
60.2	Laacher Hof	0-20 cm	92.3

5 References

1. Chemikaliengesetz, attachment d, dated July 25, 1994
2. OECD Principles of Good Laboratory Practice (GLP), dated November 26, 1997 [C(97) 186/Final]
3. [REDACTED] Method for high-performance liquid chromatographic determination of residues of the insecticide Imidacloprid in soil. Reference: MR-53/92, Method 00267 dated January 23, 1992
4. [REDACTED] Residue Analytical Method for the Determination of Residues of Imidacloprid, Hydroxy-Metabolite and Olefin-Metabolite in Nectar, Honey, Rape Flower, Rape Pollen and Bee Samples by HPLC with Electrospray MS/MS detection. Reference: MR-551/98, Method 00537 dated January 15, 1999

APPENDIX II: Plot History and Cultivation of the Plots during the Study.

Crop management before 1999 was not conducted and recorded under GLP regulations.

- Control plot: field number 711
- Variant „1997“: field number 710
- Variant „1998“: field number 702
- Variant „1998 (2x)“: field number A XII
- Variant „1999“: field number 711

Plot History

Study Plot / Year	Cropping	Pesticidal Treatments	Fertilizer Treatments
Control			
1996	Grass (<i>Lolium perenne</i>)	None	none
1997	Grass (<i>Lolium perenne</i>)	None	250 kg/ha KAS 67.5 kg/ha N
1998	Grass (<i>Lolium perenne</i>)	0.03 kg/ha Höstar [H] 1.5 L/ha Starane [H] 2 L/ha U46 M Fluid [H]	234 kg/ha KAS 63.2 kg/ha N
Variant 1999			
1996	Grass (<i>Lolium perenne</i>)	None	none
1997	Grass (<i>Lolium perenne</i>)	None	250 kg/ha KAS 67.5 kg/ha N
1998	Grass (<i>Lolium perenne</i>)	0.03 kg/ha Höstar [H] 1.5 L/ha Starane [H] 2 L/ha U46 M Fluid [H]	234 kg/ha KAS 63.2 kg/ha N
Variant 1997			
1996	Winter barley	0.5 L/dt Baytan Universal fl. [F] 0.7 L/ha Composan [H] 0.5 L/ha Folicur [F] 1.5 L/ha Matador [F] various seed dressings partly with imidacloprid	630 kg/ha KAS 170.1 kg/ha N
1997	Grass (<i>Lolium perenne</i>) & clover winter wheat (77 g imidacloprid/ha)	Akzent & Gaucho 350 FS	
1998	winter wheat	1.4 L/ha CCC 720 [H] 0.5 L/ha Enduro [I] 0.6 L/ha Metasystox R [I] 1 L/ha Pronto [F]	759 kg/ha KAS 205 kg/ha N
	winter rye	0.15 L/dt Abarit UF [F]	

[H] = herbicide and plant growth regulators, [F] = fungicide, [I] = insecticide

APPENDIX II: cont'd.

Crop management before 1999 was not conducted and recorded under GLP regulations.

- Control plot: field number 711
- Variant „1997“: field number 710
- Variant „1998“: field number 702
- Variant „1998 (2x)“: field number A XII
- Variant „1999“: field number 711

Study Plot / Year	Cropping	Pesticidal Treatments	Fertilizer Treatments
Variant 1998			
1996	Sugar beet (111 g imidacl./ha) winter wheat	TMTD & Hymexazol & imidacloprid	1004 kg/ha KAS 672 L/ha AHL 541 kg/ha N
1997	winter wheat	0.2 L/dt Abarit OF [F] 1.5 L/ha Hora Ho [H] 1.5 L/ha Tristar [H] 5.0 L/ha Roundup [H] 1 L/ha CCC 720 [H] 0.3 L/ha Bulldock [I] 0.5 L/ha Metasystox R [I] & various fungicidal spray treatments	560 kg/ha KAS 151.2 kg/ha N
	winter rye	0.2 L/dt Panoctin 35 [F] 3.0 L/ha Econal [H]	
1998	winter rye	1.0 L/ha CCC 720 [H] 0.4 L/ha Camposan extra [H] 0.5 L/ha Enduro [I] 0.8 L/ha Harvesan [F]	582 kg/ha KAS 157.2 kg/ha N
	winter barley (48.8 g imidacl./ha)	0.5 L/dt Manta plus [F,I] 0.8 L/ha Pronto [F] 0.6 L/ha Amistar [F]	
Variant 1998 (2x)			
1996	winter rye	0.2 L/dt Panoctin 35 [F] 1.5 L/ha CCC 720 [H] 1.5 L/ha Sportak [F] 1.0 L/ha Matador [F] 0.5 L/ha Bayfidan [F]	629 kg/ha KAS 170 kg/ha N
1997	winter rye summer barley	1.5 L/ha Tristar [H] 2.5 L/ha Hora Ho [H]	397 kg/ha KAS 107.2 kg/ha N
1998	Sugar beet (105.3 g Imidacl./ha)	Thiram & Imidacloprid [F&I] 3.0 L/ha Glyphos [H] 3.0 kg/ha Domino [H] 0.3 L/ha Ethosate [H] 4.0 kg/ha Goltix 70 WG 2.0 L/ha Betanal & Progress [H] 0.5 L/ha Rakobinol [H]	592.6 kg/ha KAS 160 kg/ha N
	winter wheat (76.2 g Imidacl./ha)	Gaucha FS 350 [I] 1.5 L/ha Tristar [H] 2.5 L/ha Horatto [H] 1.0 L/ha CCC 720	

[H] = herbicide and plant growth regulators, [F] = fungicide, [I] = insecticide

APPENDIX II: cont'd.

Crop management before 1999 was not conducted and recorded under GLP regulations.

- Control plot: field number 711
- Variant „1997“: field number 710
- Variant „1998“: field number 702
- Variant „1998 (2x)“: field number A XII
- Variant „1999“: field number 711

1999 Treatments

Study Plot / Year	Cropping	Pesticidal Treatments	Fertilizer Treatments
Control	Field No. 711		
29 March	Grass (<i>Lolium perenne</i>)	4 L/ha Glyphos [H]	
12 May	summer rape	TMTD [F]	
14 May	summer rape	2.5 L/ha Butisan S [H]	
20 May	summer rape		190 kg/ha KAS 51.3 kg/ha N
1 June	summer rape	1.0 L/ha MesuroI SC 500 [I]	
28 June	summer rape	1.2 L/ha Uden [I]	
Variant 1999			
29 March	Grass (<i>Lolium perenne</i>)	4 L/ha Glyphos [H]	
12 May	summer rape [33.5 g imidacI./ha]	TMTD [F] 25 mL/dt Poncho FS 500 [I]	
14 May	summer rape	2.5 L/ha Butisan S [H]	
20 May	summer rape		190 kg/ha KAS 51.3 kg/ha N
1 June	summer rape	1.0 L/ha MesuroI SC 500 [I]	
28 June	summer rape	1.2 L/ha Uden [I]	
Variant 1997			
12 March	winter rye		221 kg/ha KAS 59.7 kg/ha N
29 March	winter rye	4 L/ha Glyphos [H]	
23 April	uncropped [50 g imidacloprid/ha]	71.5 g Gaucho WS 70 spray	
12 May	summer rape	TMTD [F]	
14 May	summer rape	2.5 L/ha Butisan S [H]	
20 May	summer rape		190 kg/ha KAS 51.3 kg/ha N
1 June	summer rape	1.0 L/ha MesuroI SC 500 [I]	
28 June	summer rape	1.2 L/ha Uden [I]	

[H] = herbicide and plant growth regulators, [F] = fungicide, [I] = insecticide

APPENDIX II: cont'd.

Crop management before 1999 was not conducted and recorded under GLP regulations.

- Control plot: field number 711
- Variant „1997“: field number 710
- Variant „1998“: field number 702
- Variant „1998 (2x)“: field number A XII
- Variant „1999“: field number 711

Study Plot / Year	Cropping	Pesticidal Treatments	Fertilizer Treatments
Variant 1998			
12 March	Winter rye		221 kg/ha KAS 59.7 kg/ha N
29 March	Winter rye	4 L/ha Glyphos [H]	
12 May	summer rape	TMTD [F]	
14 May	summer rape	2.5 L/ha Butisan S [H]	
20 May	summer rape		190 kg/ha KAS 51.3 kg/ha N
1 June	summer rape	1.0 L/ha Mesurool SC 500 [I]	
28 June	summer rape	1.2 L/ha Uden [I]	
Variant 1998 (2x)			
12 March	Winter wheat		221 kg/ha KAS 59.7 kg/ha N
29 March	Winter wheat	5 L/ha Glyphos [H]	
20 April	uncropped		182 kg/ha KAS 49.1 kg/ha N
12 May	summer rape	TMTD [F]	
14 May	summer rape	2.5 L/ha Butisan S [H]	
20 May	summer rape		190 kg/ha KAS 51.3 kg/ha N
1 June	summer rape	1.0 L/ha Mesurool SC 500 [I] 1.2 L/ha Uden [I]	

[H] = herbicide and plant growth regulators, [F] = fungicide, [I] = insecticide

Dit document is geen eigendom van het Ctgb en wordt beschermd op grond van de wettelijke verplichting tot openbaarmaking. Op dit document kunnen rechten van de auteursrechten, in het bijzonder intellectuele eigendomsrechten en/of auteursrechten. Voorts kan dit document onder een regeling omtrent gegevensbescherming vallen. Het is niet toegestaan dit document of de inhoud hiervan te kopiëren of te verspreiden zonder de toestemming van de rechthebbende van dit document. Het is niet toegestaan dit document te verspreiden of te verspreiden van de rechten van de rechthebbende.

This document is not the property of the Ctgb and is only provided based on mandatory freedom of information requirements. The document may be subject to rights such as intellectual property and copy rights of third parties. Consequently, any publication, distribution, reproduction and/or publishing and any commercial exploitation and use of this document or its contents without the permission of the owner of this document are prohibited and violate the rights of its owner.

APPENDIX III: Analytical Report for Biological Samples.

Bayer AG
Crop Protection Development
Institute for Metabolism Research
and Residue Analysis
D-51368 Leverkusen

September 28, 1999
Report No.: MR-518/99
Page 31 of 43

STUDY TITLE

Residue Levels of Imidacloprid and Imidacloprid Metabolites in Nectar, Blossoms and Pollen of Summer Rape Cultivated on Soils with Different Imidacloprid Residue Levels and Effects of These Residues on Foraging Honeybees

Test Location: farmland "Laacher Hof"

Author

[Redacted]

Testing Facility

Bayer AG
PF-E/MR, Building 6610
51368 Leverkusen, Germany

Study Completion Date

September 28, 1999

Study Number

E 370 1548-8

Dit document is geen eigendom van het Ctgb en wordt beschikbaar gemaakt op grond van een wettelijk verplichting tot openbaarmaking. Op dit document kunnen rechten van derden rusten, waaronder intellectuele eigendomsrechten. Voorts kan dit document onder een regeling omtrent gegevensbescherming vallen. Het gebruik van dit document is de inhoud hiervan te onder de toestemming van de rechthebbende van dit document kan derhalve verboden zijn en een inbreuk opleveren van de rechten van de rechthebbende.

This document is not the property of the Ctgb and is only provided based on mandatory freedom of information requirements. The document may be subject to rights such as intellectual property and copy rights of third parties. Furthermore, this document may fall under a regulatory data protection regime. Consequently, any publication, distribution, reproduction and/or publishing and any commercial exploitation and use of this document or its contents without the permission of the owner of this document may therefore be prohibited and violate the rights of its owner.

TABLE OF CONTENT

	Page No.
STUDY TITLE	31
TABLE OF CONTENT	32
1 INTRODUCTION	33
2 TIME SCHEDULE	34
3 RESULTS OF NECTAR, POLLEN, FLOWER, BEE AND GREEN	
MATERIAL SAMPLES :	33
3.1 Nectar Samples:	33
3.2 Pollen Samples:	34
3.3 Flower Samples:	34
3.4 Bee Samples:	35
3.5 Green Material Samples:	35
4 EXPERIMENTAL	36
4.1 Reference Substances	36
4.2 Residue Analytical Methodology	37
4.2.1 Extraction and Sample Clean-up	37
4.2.2 ChemElut® Column Clean-up	37
4.2.3 Silica Gel Column Clean-up	38
4.3 HPLC-MS/MS determination of Imidacloprid and Metabolites	39
4.3.1 Measuring equipment and HPLC conditions:	39
4.3.2 MS/MS Detection	40

Dit document is geen eigendom van het Ctgb en wordt niet beschermd door intellectuele eigendomsrechten en/of auteursrechten. Voorts kan dit document of de inhoud hiervan zonder de toestemming van de rechthebbende van dit document of de rechten van deze rechthebbende.

This document is not the property of the Ctgb and only provided based on mandatory freedom of information requirements. The document may be subject to rights such as intellectual property and copy rights of third parties. Furthermore, this document may fall under a regulatory data protection regime. Consequently, any publication, distribution, reproduction and/or publishing and any commercial exploitation and use of this document or its contents without the permission of the owner of this document may therefore be prohibited and violate the rights of its owner.

1 INTRODUCTION

Summer rape samples of the Germany trial station "Laacher Hof" were analysed for residues of Imidacloprid and its Olefin- and Hydroxy Metabolites. The results are tabulated in the table below. Extraction, sample clean up and determination of Imidacloprid, Hydroxy- and Olefin-Metabolite by HPLC-MS/MS were performed according to method 00537/E001 (MR-568/99). The limit of quantitation was 0.005 mg/kg for Imidacloprid and the Hydroxy-Metabolite and 0.01 mg/kg for the Olefin-Metabolite. The limit of detection was 0.0015 mg/kg for Imidacloprid and the Hydroxy-Metabolite and 0.003 mg/kg for the Olefin-Metabolite.

2 TIME SCHEDULE

The experimental work was performed during the following time period.

Signature of Study Protocol: March 22, 1999
 Start of Experimental Phase: August 30, 1999
 End of Experimental Phase: September 21, 1999
 Completion of Report: September 28, 1999

3 RESULTS OF NECTAR, POLLEN, FLOWER, BEE AND GREEN

MATERIAL SAMPLES :

3.1 Nectar Samples:

Sample Name	Sample description	Sample weight [g]	Hydroxy-NTN [mg/kg]	Olefin-NTN [mg/kg]	Imidacloprid [mg/kg]
E15488K001	Nectar	0.37	n.d.	n.d.	n.d.
E15488E97001	Nectar	0.57	n.d.	n.d.	n.d.
E15488E98001	Nectar	0.8	n.d.	n.d.	n.d.
E15488E99001	Nectar	0.5	n.d.	n.d.	< LOQ
E15488D98001	Nectar	0.5	n.d.	n.d.	n.d.

Limit of quantitation: 0.005 mg/kg for Imidacloprid and Hydroxy-Metabolite, 0.01 mg/kg for the Olefin-Metabolite; < 0.005 and < 0.010 = Residues below the limit of quantitation

Limit of detection: 0.0015 mg/kg for Imidacloprid and Hydroxy-Metabolite, 0.003 mg/kg for the Olefin-Metabolite; n.d.: Residues below the limit of detection

3.2 Pollen Samples:

Sample Name	Sample description	Sample weight [g]	Hydroxy-NTN [mg/kg]	Olefin-NTN [mg/kg]	Imidacloprid [mg/kg]
E15488K003	Pollen	1.35	n.d.	n.d.	n.d.
E15488E97003	Pollen	1.47	n.d.	n.d.	n.d.
E15488E98003	Pollen	1.32	n.d.	n.d.	< LOQ
E15488E99003	Pollen	0.57	n.d.	n.d.	n.d.
E15488D98003	Pollen	0.4	n.d.	n.d.	< LOQ

Limit of quantitation: 0.005 mg/kg for Imidacloprid and Hydroxy-Metabolite, 0.01 mg/kg for the Olefin-Metabolite; < 0.005 and < 0.010 = Residues below the limit of quantitation.

Limit of detection: 0.0015 mg/kg for Imidacloprid and Hydroxy-Metabolite, 0.003 mg/kg for the Olefin-Metabolite; n.d.: Residues below the limit of detection.

3.3 Flower Samples:

Sample Name	Sample description	Sample weight [g]	Hydroxy-NTN [mg/kg]	Olefin-NTN [mg/kg]	Imidacloprid [mg/kg]
E15488K002	Rape Flower	5.6	n.d.	n.d.	n.d.
E15488E97002	Rape Flower	7.2	n.d.	n.d.	n.d.
E15488E98002	Rape Flower	8.0	n.d.	n.d.	n.d.
E15488E99002	Rape Flower	5.7	n.d.	n.d.	n.d.
E15488D98002	Rape Flower	7.4	n.d.	n.d.	n.d.

Limit of quantitation: 0.005 mg/kg for Imidacloprid and Hydroxy-Metabolite, 0.01 mg/kg for the Olefin-Metabolite; < 0.005 and < 0.010 = Residues below the limit of quantitation.

Limit of detection: 0.0015 mg/kg for Imidacloprid and Hydroxy-Metabolite, 0.003 mg/kg for the Olefin-Metabolite; n.d.: Residues below the limit of detection.

Dit document is een eigendom van het Ctgb en wordt beschikbaar gemaakt op grond van een wettelijke verplichting tot openbaarmaking. Op dit document kan rechten van auteursrecht, intellectuele eigendomsrechten en/of auteursrecht, waaronder intellectuele eigendomsrechten en/of auteursrechten, voortvloeien en de afzender aanvaardt de aansprakelijkheid daarvan. Het is niet toegestaan dit document te kopiëren, te verspreiden of openbaar te maken zonder de toestemming van de afzender. Het gebruik van dit document is beperkt tot de afzender en kan vertrouwelijk of anderszins beschermd zijn. Het is niet toegestaan dit document te kopiëren, te verspreiden of openbaar te maken zonder de toestemming van de afzender. Het gebruik van dit document is beperkt tot de afzender en kan vertrouwelijk of anderszins beschermd zijn.

This document is not the property of the Ctgb and only provided based on mandatory freedom of information requirements. Consequently, any publication, distribution, reproduction and use of this document or its contents without the permission of the owner of this document may therefore be prohibited and violate the rights of its owner. Furthermore, this document may fall under a regulatory data protection regime.

3.4 Bee Samples:

Sample Name	Sample description	Sample weight [g]	Hydroxy-NTN [mg/kg]	Olefin-NTN [mg/kg]	Imidacloprid [mg/kg]
E15488K004	Bees of Flowers	12.1	n.d.	n.d.	n.d.
E15488E97004	Bees of Flowers	8.4	n.d.	n.d.	n.d.
E15488E98004	Bees of Flowers	11.3	n.d.	n.d.	n.d.
E15488E99004	Bees of Flowers	9.9	n.d.	n.d.	n.d.
E15488D98004	Bees of Flowers	8.3	n.d.	n.d.	n.d.

Limit of quantitation: 0.005 mg/kg for Imidacloprid and Hydroxy-Metabolite, 0.01 mg/kg for the Olefin-Metabolite; < 0.005 and < 0.010 = Residues below the limit of quantitation.

Limit of detection: 0.0015 mg/kg for Imidacloprid and Hydroxy-Metabolite, 0.003 mg/kg for the Olefin-Metabolite; n.d.: Residues below the limit of detection.

3.5 Green Material Samples:

Sample Name	Sample description	Sample weight [g]	Hydroxy-NTN [mg/kg]	Olefin-NTN [mg/kg]	Imidacloprid [mg/kg]
E15488K006	Green Material	28	n.d.	n.d.	n.d.
E15488E97006	Green Material	69	n.d.	n.d.	< LOQ
E15488E98006	Green Material	67	n.d.	n.d.	< LOQ
E15488E99006	Green Material	69	< LOQ	n.d.	< LOQ
E15488D98006	Green Material	63	< LOQ	n.d.	< LOQ

Limit of quantitation: 0.005 mg/kg for Imidacloprid and Hydroxy-Metabolite, 0.01 mg/kg for the Olefin-Metabolite; < 0.005 and < 0.010 = Residues below the limit of quantitation.


Limit of detection: 0.0015 mg/kg for Imidacloprid and Hydroxy-Metabolite, 0.003 mg/kg for the Olefin-Metabolite; n.d.: Residues below the limit of detection.

4 EXPERIMENTAL

4.1 Reference Substances

Imidacloprid

Structural formula:



Empirical formula:

 $C_9H_{10}ClN_5O_2$

Molecular weight:

255.7 g/mole

Certificate of Analysis:

M00680, 03/13/98

Certified Assay:


99.4 %

Expiry Date:

March 2000

Hydroxy-Imidacloprid (WAK 4103)

Structural formula:



Empirical formula:

 $C_9H_{10}ClN_5O_4$

Molecular weight:

271.7 g/mole

Certificate of Analysis:

930323ELB03, 06/07/95

Certified Assay:


99.4 %

Expiry Date:

June 2000

Olefin-Imidacloprid (NTN 35884)

Structural formula:



Empirical formula:

 $C_9H_8ClN_5O_2$

Molecular weight:

253.6 g/mole

Certificate of Analysis:

M00804, 07/22/98

Certified Assay:

98 %

Expiry Date:

June 2000

4.2 Residue Analytical Methodology

4.2.1 Extraction and Sample Clean-up

1. Place for e.g. 2.0 g of the sample material in a 150-ml beaker.
Add 30 ml of methanol/water (3/1, v/v) and allow the sample to soak for 30 min.
2. Blend the sample using an ultra-turrax blender (or equivalent) for approximately 1 min.
3. Vacuum filter the suspension through 2.5 g of Celite filter aid using Schwarzband filter paper supported on a Büchner funnel into a 250-ml vacuum filter flask.
4. Wash the filtered solids with a total of 30 ml of methanol/water (3/1, v/v). Press residual solvent from the solids using rubber damming. Discard the filtered solids.
5. Transfer the filtrate to a 100-ml graduated cylinder. Determine the total volume of the extracts. Mix the solution well, and transfer the half (e.g. 1.0 g sample equivalent) to a 250-ml brown glass round-bottomed flask.
6. Concentrate the aliquot to an aqueous remainder of 5 to 10 ml using a rotary evaporator with a max. bath temperature of 50 °C.

4.2.2 ChemElut[®] Column Clean-up

1. Add 5 to 10 ml water to the aqueous solution from 4.2.1 step 6 to bring the total volume of the extracts to approx. 20 ml.
2. Place the aqueous solution on the top of the ChemElut[®] CE 1020 (20 ml volume) column fitted with a disposable stainless steel needle and wait for approx. 15 minutes to achieve an uniform distribution of the liquid on the column.
3. Elute the residues from the column with 140 ml of CH₂Cl₂. Collect the eluate in a 250-ml brown glass round-bottomed flask.
4. Evaporate the eluate from step 3 to dryness using a vacuum rotary evaporator and a max. bath temperature of 40 °C.

4.2.3 Silica Gel Column Clean-up

1. Dissolve the residues from 4.2.2 step 4 in 2 ml of toluene/ethyl acetate (85/15, v/v).
2. Apply the organic solution from step 1 onto a 0.5 g (3 ml) silica gel (SiOH) column (e.g. Varian).
3. Allow the solution to pass through the column at a flow rate of 1 ml/min.
4. Rinse the 250-ml brown glass round-bottomed flask with 10 ml of toluene/ethyl acetate (70/30, v/v) and apply the solution onto the column, too.
5. Elute the residues with 5 ml of acetonitrile at a flow rate of 1 ml/min. Collect the eluate in a 25-ml brown glass pear-shaped flask.
6. Evaporate the eluate from step 5 to dryness using a vacuum rotary evaporator and a max. bath temperature of 40 °C. Dissolve the residues in e.g. 1.00 ml of acetonitrile/water (2/8, v/v) and determine the residues with HPLC-MS/MS.

NOTE

1. The volumes to be used for flushing the column with toluene/ethyl acetate and for elution with acetonitrile must be newly determined for each batch of SiOH-column!

2. The flow rate should not be too high, since otherwise losses of the residues in may occur with recoveries below 70 % and the clean-up is less effective.

3. The Hydroxy-Metabolite may be converted to the Olefin-Metabolite (especially under acidic conditions).

4. The Olefin-Metabolite is degraded by light (ca. 50% in one day at natural daylight). Therefore, all solutions containing the Olefin-Metabolite must be protected from light and stored in a cool and dark place.

4.3 HPLC-MS/MS determination of Imidacloprid and Metabolites

4.3.1 Measuring equipment and HPLC conditions:

Instrument: HP 1100
 Injector: HP 1100
 Column: Phenomenex, Luna C18 (2), 5 µm, 15 cm, 0.46 cm i.d.
 or equivalent
 Injection Volume: 50 µl
 Oven temperature: 40 °C
 Mobile Phase: A: Water/ACN (90/10, v/v)+ 0.1 ml acetic acid per litre
 B: Acetonitrile + 0.1 ml acetic acid per litre

Time Table	0 min	11.1 % B
	10 min	11.1 % B
	10.1 min	90 % B
	15 min	90 % B
	15.1 min	11.1 % B
	19 min	11.1 % B

Stoptime: 19 min
 Flow (Column): 1.0 ml/min
 Flow (into MS): 0.15 ml/min
 Retention Time: Olefin-Metabolite: approx. 4.6 min
 Hydroxy-Metabolite: approx. 5.5 min
 Imidacloprid: approx. 9.1 min

NOTE: Conditions may be adapted for other HPLC-MS/MS systems.

Dit document is geen eigendom van het Ctgb en wordt beschikbaar gemaakt op grond van een wettelijke verplichting tot openbaarmaking. Op dit document kunnen rechten van auteursrechten, intellectuele eigendomsrechten of auteursrechten. Publicatie, verspreiding, vermenigvuldiging, commerciële exploitatie en gebruik van dit document behouden zijn en een inbreuk op deze rechten van deze rechthebbende. This document is not the property of the Ctgb and is provided based on mandatory freedom of information requirements. The document may be subject to rights such as intellectual property and copy rights of third parties. Furthermore, this document may fall under a regulatory data protection regime. Consequently, any publication, distribution, reproduction and/or publishing and any commercial exploitation and use of this document or its contents without the permission of the owner of this document may therefore be prohibited and violate the rights of its owner.

4.3.2 MS/MS-Detection

The experiments were performed on a triple-quadrupole mass spectrometer system, fitted with an electrospray interface operated in the positive ion mode under MRM conditions.

The mass spectrometer was tuned by infusing a standard solution of 0.5 mg/l Imidacloprid and its metabolites (dissolved in water/acetonitrile 8/2 + 0.1 ml acetic acid per l) at a flow rate of 10-20 µl/min. Mass axis calibration was done by infusing a polypropylene glycol 3000 solution. Unit mass resolution was established and maintained in each mass resolving quadrupole by maintaining a full width at half-maximum of between 0.8 and 1.0 DA. After tuning and calibration, optimal collision-activated dissociation (CAD) conditions for fragmentation of Imidacloprid and its metabolites were determined. These experiments were performed with nitrogen as collision gas with a collision offset of -19 eV for Imidacloprid, -21 eV for the Hydroxy-Metabolite and -13 eV for the Olefin-Metabolite and at an approximate collision gas thickness of 1.46×10^{15} atoms/cm². Nebulizer gas is set at 1.48 l/min, curtain gas is set at 1.44 l/min collision gas is set at 0.87 l/min and turbo gas is set at 6.0 l/min.

Detector: Triple Quadrupole MS/MS Mass Spectrometer, e.g. Perkin-Elmer Sciex Instruments API 300, Apple™ Macintosh System® 8.1

Interface: Electrospray, Turbo Ion Spray
 Potential: +4400 V
 Temperature: 400 °C
 Nebulizer Gas: Nitrogen 5.0 (99.999 % purity), 1.48 l/min
 Curtain Gas: Nitrogen 5.0 (99.999 % purity), 1.44 l/min
 Turbo Gas: Nitrogen 5.0 (99.999 % purity), 6.0 l/min

Scan Type: MRM (Multiple Reaction Monitoring Mode)

Polarity: Positive

Collision Gas: Nitrogen 5.0 (99.999 % purity), 0.87 l/min


Mass spectrometer operating parameters.

Compound	Precursor Ion Q1 Mass (amu)	Product Ion Q3 Mass (amu)	Dwell Time (msec)	Collision Energy (eV)
Olefin-Metabolite (37)	256#	238	250	-13
Olefin-Metabolite (35)	254	236	250	-13
Hydroxy-Metabolite (37)	274#	191	250	-21
Hydroxy-Metabolite (35)	272	191	250	-21
Imidacloprid (37)	258#	211	500	-19
Imidacloprid (35)	256	209	500	-19

#: The Cl 37 isotope of all substances was detected to build the isotopes ratio

NOTE: Different MS/MS-instruments or instrument parameters may result in different ion transitions and different relative intensities.

Appendix IV: Copy of the GLP Certificate




baarmaking.
ing van de

ents.
his document or its contents
its of its owner.

Dit document is geen eigend
Op dit docume
Publicatie, verspreiding
rechthe

Consequently, any p
without

Appendix IV: cont`d.




Dit document is geen eigendom van de afzender.
Op dit document kunnen auteursrechten of andere rechten van derden van toepassing zijn.
Publicatie, verspreiding, vermenigvuldiging, verspreiding of anderszins openbaar maken van dit document kan strafbaar zijn.
Voorts kan het gebruik van dit document strafbaar zijn.
This document is not the property of the Ctgb or the applicant.
The document may be subject to rights of third parties.
Consequently, any publication, distribution, reproduction and/or public use of this document without the permission of the owner of this document may be liable to legal action.

baarmaking.
ing van de
ements.
of this document or its contents
rights of its owner.

Appendix V: Quality Assurance Statement

Referat GLP

Quality Assurance Statement



haarmaking.
ing van de

ntents

Dit d